Publications of William Trischuk

April 1, 2013

** Indicates papers where I played a leading role in the analysis, manuscript preparation or review.

Papers submitted for publication

- 1. G. Aad et al. [ATLAS Collaboration], "Measurement with the ATLAS detector of multiparticle azimuthal correlations in p + Pb collisions at $\sqrt{s_{NN}}$ =5.02 TeV," arXiv:1303.2084 [hep-ex].
- 2. G. Aad et al. [ATLAS Collaboration], "Search for third generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," arXiv:1303.0526 [hep-ex].
- 3. ** G. Aad *et al.* [ATLAS Collaboration], "Characterisation and mitigation of beaminduced backgrounds observed in the ATLAS detector during the 2011 proton-proton run," arXiv:1303.0223 [hep-ex].
- 4. G. Aad et al. [ATLAS Collaboration], "Search for WH production with a light Higgs boson decaying to prompt electron-jets in proton-proton collisions at \sqrt{s} =7 TeV with the ATLAS detector," arXiv:1302.4403 [hep-ex].
- 5. ** G. Aad *et al.* [ATLAS Collaboration], "Improved luminosity determination in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC," arXiv:1302.4393 [hep-ex].
- 6. G. Aad et al. [ATLAS Collaboration], "Search for a light charged Higgs boson in the decay channel $H^+ \to c\bar{s}$ in ttbar events using pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," arXiv:1302.3694 [hep-ex].
- 7. G. Aad et al. [The ATLAS Collaboration], "Measurement of the cross-section for W boson production in association with b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," arXiv:1302.2929 [hep-ex].
- 8. G. Aad et al. [The ATLAS Collaboration], "Measurement of kT splitting scales in $W \to l\nu$ events at \sqrt{s} =7 TeV with the ATLAS detector," arXiv:1302.1415 [hep-ex].
- 9. G. Aad et al. [The ATLAS Collaboration], "Measurements of Wgamma and Zgamma production in pp collisions at \sqrt{s} = 7 TeV with the ATLAS detector at the LHC," arXiv:1302.1283 [hep-ex].
- 10. G. Aad et al. [ATLAS Collaboration], "Search for long-lived, multi-charged particles in pp collisions at \sqrt{s} =7 TeV using the ATLAS detector," arXiv:1301.5272 [hep-ex].
- 11. G. Aad *et al.* [ATLAS Collaboration], "Search for single b^* -quark production with the ATLAS detector at $\sqrt{s} = 7$ TeV," arXiv:1301.1583 [hep-ex].

- 12. G. Aad *et al.* [ATLAS Collaboration], "Observation of Associated Near-side and Away-side Long-range Correlations in $\sqrt{s_{NN}} = 5.02$ TeV Proton-lead Collisions with the ATLAS Detector," arXiv:1212.5198 [hep-ex].
- 13. G. Aad *et al.* [ATLAS Collaboration], "Search for charged Higgs bosons through the violation of lepton universality in $t\bar{t}$ events using pp collision data at $\sqrt{s}=7$ TeV with the ATLAS experiment," arXiv:1212.3572 [hep-ex].
- 14. G. Aad et al. [ATLAS Collaboration], "Search for a heavy narrow resonance decaying to $e\mu$, $e\tau$, or $\mu\tau$ with the ATLAS detector in $\sqrt{s}=7$ TeV pp collisions at the LHC," arXiv:1212.1272 [hep-ex].
- 15. G. Aad *et al.* [ATLAS Collaboration], "Measurement of Upsilon production in 7 TeV *pp* collisions at ATLAS," arXiv:1211.7255 [hep-ex].
- 16. G. Aad et al. [ATLAS Collaboration], "Measurement of the $t\bar{t}$ production cross section in the τ +jets channel using the ATLAS detector," arXiv:1211.7205 [hep-ex].
- 17. G. Aad et al. [ATLAS Collaboration], "Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at \sqrt{s} =7 TeV with the ATLAS detector," arXiv:1211.6956 [hep-ex].
- 18. G. Aad *et al.* [ATLAS Collaboration], "Measurement of angular correlations in Drell-Yan lepton pairs to probe Z/γ^* boson transverse momentum at \sqrt{s} =7 TeV with the ATLAS detector," arXiv:1211.6899 [hep-ex].
- 19. G. Aad et al. [ATLAS Collaboration], "Measurement of W^+W^- production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector and limits on anomalous WWZ and $WW\gamma$ couplings," arXiv:1210.2979 [hep-ex].
- 20. G. Aad *et al.* [ATLAS Collaboration], "Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector," arXiv:1210.4491 [hep-ex].
- 21. G. Aad et al. [ATLAS Collaboration], "Measurement of charged-particle event shape variables in $\sqrt{s} = 7$ TeV proton-proton interactions with the ATLAS detector," arXiv:1207.6915 [hep-ex].
- 22. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the Λ_b lifetime and mass in the ATLAS experiment," arXiv:1207.2284 [hep-ex].

Papers published in refereed journals

1. ** G. Aad *et al.* [ATLAS Collaboration], "A particle consistent with the Higgs Boson observed with the ATLAS Detector at the Large Hadron Collider," Science **338**, 1576 (2012).

- 2. G. Aad et al. [ATLAS Collaboration], "Multi-channel search for squarks and gluinos in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector," Eur. Phys. J. C 73, 2362 (2013).
- 3. G. Aad et al. [ATLAS Collaboration], "A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **719**, 299 (2013).
- 4. G. Aad *et al.* [ATLAS Collaboration], "Search for new phenomena in events with three charged leptons at a center-of-mass energy of 7 TeV with the ATLAS detector," Phys. Rev. D **87**, 052002 (2013).
- 5. G. Aad et al. [ATLAS Collaboration], "Measurement of ZZ production in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous ZZZ and $ZZ\gamma$ couplings with the ATLAS detector," JHEP **1303**, 128 (2013).
- 6. G. Aad et al. [ATLAS Collaboration], "Search for resonances decaying into top-quark pairs using fully hadronic decays in pp collisions with ATLAS at $\sqrt{s} = 7$ TeV," JHEP **1301**, 116 (2013).
- 7. G. Aad et al. [ATLAS Collaboration], "Measurement of isolated-photon pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1301**, 086 (2013).
- 8. G. Aad *et al.* [ATLAS Collaboration], "Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Phys. Lett. B **720**, 277 (2013).
- 9. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry in events with photons, bottom quarks, and missing transverse momentum in proton-proton collisions at a centre-of-mass energy of 7 TeV with the ATLAS detector," Phys. Lett. B **719**, 261 (2013).
- 10. G. Aad et al. [ATLAS Collaboration], "Search for contact interactions and large extra dimensions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 87, 015010 (2013).
- 11. G. Aad *et al.* [ATLAS Collaboration], "Search for Extra Dimensions in diphoton events using proton-proton collisions recorded at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC," New J. Phys. **15**, 043007 (2013).
- 12. G. Aad *et al.* [ATLAS Collaboration], "Search for long-lived, heavy particles in final states with a muon and multi-track displaced vertex in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **719**, 280 (2013).
- 13. G. Aad *et al.* [ATLAS Collaboration], "A search for high-mass resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector," Phys. Lett. B **719**, 242 (2013).
- 14. G. Aad *et al.* [ATLAS Collaboration], "Measurement of Z boson Production in Pb + Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector," Phys. Rev. Lett. **110**, 022301 (2013).

- 15. G. Aad *et al.* [ATLAS Collaboration], "Jet energy resolution in proton-proton collisions at $\sqrt{s} = 7$ TeV recorded in 2010 with the ATLAS detector," Eur. Phys. J. C **73**, 2306 (2013).
- 16. G. Aad et al. [ATLAS Collaboration], "Search for pair production of heavy top-like quarks decaying to a high-pT W boson and a b quark in the lepton plus jets final state at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **718**, 1284 (2013).
- 17. G. Aad *et al.* [ATLAS Collaboration], "Search for doubly-charged Higgs bosons in like-sign dilepton final states at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C **72**, 2244 (2012).
- 18. G. Aad *et al.* [ATLAS Collaboration], "Search for pair-produced massive coloured scalars in four-jet final states with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C **73**, 2263 (2013).
- 19. G. Aad *et al.* [ATLAS Collaboration], "Search for pair production of massive particles decaying into three quarks with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC," JHEP **1212**, 086 (2012).
- 20. G. Aad *et al.* [ATLAS Collaboration], "Search for anomalous production of prompt likesign lepton pairs at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1212**, 007 (2012).
- 21. G. Aad *et al.* [ATLAS Collaboration], "Search for R-parity-violating supersymmetry in events with four or more leptons in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector," JHEP **1212**, 124 (2012).
- 22. G. Aad *et al.* [ATLAS Collaboration], "Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1301**, 131 (2013).
- 23. G. Aad *et al.* [ATLAS Collaboration], "ATLAS search for new phenomena in dijet mass and angular distributions using pp collisions at $\sqrt{s} = 7$ TeV," JHEP **1301**, 029 (2013).
- 24. G. Aad *et al.* [ATLAS Collaboration], "Search for Supersymmetry in Events with Large Missing Transverse Momentum, Jets, and at Least One Tau Lepton in 7 TeV Proton-Proton Collision Data with the ATLAS Detector," Eur. Phys. J. C 72, 2215 (2012).
- 25. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the flavour composition of dijet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C 73, 2301 (2013).
- 26. G. Aad et al. [ATLAS Collaboration], "Search for displaced muonic lepton jets from light Higgs boson decay in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **721**, 32 (2013).
- 27. G. Aad et al. [ATLAS Collaboration], "Search for resonant top plus jet production in $t\bar{t}$ + jets events with the ATLAS detector in pp collisions at $\sqrt{s}=7$ TeV," Phys. Rev. D 86, 091103 (2012).

- 28. G. Aad *et al.* [ATLAS Collaboration], "Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. Lett. **110**, 011802 (2013).
- 29. G. Aad et al. [ATLAS Collaboration], "ATLAS search for a heavy gauge boson decaying to a charged lepton and a neutrino in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C 72, 2241 (2012).
- 30. G. Aad *et al.* [ATLAS Collaboration], "Search for a heavy top-quark partner in final states with two leptons with the ATLAS detector at the LHC," JHEP **1211**, 094 (2012).
- 31. G. Aad *et al.* [ATLAS Collaboration], "Search for high-mass resonances decaying to dilepton final states in pp collisions at $\sqrt{s} = 7$ -TeV with the ATLAS detector," JHEP **1211**, 138 (2012).
- 32. G. Aad *et al.* [ATLAS Collaboration], "Search for light top squark pair production in final states with leptons and b^- jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions," Phys. Lett. B **720**, 13 (2013).
- 33. J. Albert *et al.* [ATLAS IBL Collaboration], "Prototype ATLAS IBL Modules using the FE-I4A Front-End Readout Chip," JINST 7, P11010 (2012).
- 34. G. Aad *et al.* [ATLAS Collaboration], "Search for diphoton events with large missing transverse momentum in 7 TeV proton-proton collision data with the ATLAS detector," Phys. Lett. B **718**, 411 (2012).
- 35. G. Aad *et al.* [ATLAS Collaboration], "Measurements of the pseudorapidity dependence of the total transverse energy in proton-proton collisions at $\sqrt{s} = 7$ TeV with ATLAS," JHEP **1211**, 033 (2012).
- 36. G. Aad *et al.* [ATLAS Collaboration], "Further search for supersymmetry at $\sqrt{s} = 7$ TeV in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector," Phys. Rev. D **86**, 092002 (2012).
- 37. G. Aad et al. [ATLAS Collaboration], "Search for light scalar top quark pair production in final states with two leptons with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions," Eur. Phys. J. C 72, 2237 (2012).
- 38. G. Aad *et al.* [ATLAS Collaboration], "Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector," Phys. Lett. B **718**, 841 (2013).
- 39. G. Aad et al. [ATLAS Collaboration], "Search for direct slepton and gaugino production in final states with two leptons and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Phys. Lett. B **718**, 879 (2013).
- 40. G. Aad et al. [ATLAS Collaboration], "Search for new phenomena in the WW to $\ell\nu\ell$ ' ν ' final state in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **718**, 860 (2013).

- 41. G. Aad *et al.* [ATLAS Collaboration], "Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions using 4.7 fb^{-1} of ATLAS data," Phys. Rev. Lett. **109**, 211803 (2012).
- 42. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the jet radius and transverse momentum dependence of inclusive jet suppression in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector," Phys. Lett. B **719**, 220 (2013).
- 43. G. Aad *et al.* [ATLAS Collaboration], "Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. Lett. **109**, 211802 (2012).
- 44. G. Aad *et al.* [ATLAS Collaboration], "Measurement of WZ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C **72**, 2173 (2012).
- 45. G. Aad *et al.* [ATLAS Collaboration], "Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb⁻¹ of $\sqrt{s} = 7$ TeV proton-proton collision data," Phys. Rev. D **87**, 012008 (2013).
- 46. G. Aad et al. [ATLAS Collaboration], "Time-dependent angular analysis of the decay $B_s \to J/\psi \phi$ and extraction of Delta Γ_s and the CP-violating weak phase ϕ_s by ATLAS," JHEP **1212**, 072 (2012).
- 47. G. Aad et al. [ATLAS Collaboration], "Underlying event characteristics and their dependence on jet size of charged-particle jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 86, 072004 (2012).
- 48. ** G. Aad *et al.* [ATLAS Collaboration], "Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC," Phys. Lett. B **716**, 1 (2012).
- 49. G. Aad *et al.* [ATLAS Collaboration], "Search for magnetic monopoles in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector," Phys. Rev. Lett. **109**, 261803 (2012).
- 50. G. Aad et al. [ATLAS Collaboration], "Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C 73, 2261 (2013).
- 51. G. Aad *et al.* [ATLAS Collaboration], "Search for top and bottom squarks from gluino pair production in final states with missing transverse energy and at least three *b*-jets with the ATLAS detector," Eur. Phys. J. C **72**, 2174 (2012).
- 52. G. Aad *et al.* [ATLAS Collaboration], "A search for $t\bar{t}$ resonances in lepton+jets events with highly boosted top quarks collected in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1209**, 041 (2012).
- 53. G. Aad *et al.* [ATLAS Collaboration], "Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D **86**, 032003 (2012).

- 54. G. Aad *et al.* [ATLAS Collaboration], "Search for the Standard Model Higgs boson produced in association with a vector boson and decaying to a *b*-quark pair with the ATLAS detector," Phys. Lett. B **718**, 369 (2012).
- 55. G. Aad *et al.* [ATLAS Collaboration], "Search for the Higgs boson in the $H \to WW \to$ lnujj decay channel at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **718**, 391 (2012).
- 56. G. Aad et al. [ATLAS Collaboration], "Search for the Standard Model Higgs boson in the H to $\tau^+\tau^-$ decay mode in $\sqrt{s}=7$ TeV pp collisions with ATLAS," JHEP **1209**, 070 (2012).
- 57. G. Aad *et al.* [ATLAS Collaboration], "ATLAS measurements of the properties of jets for boosted particle searches," Phys. Rev. D **86**, 072006 (2012).
- 58. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the b-hadron production cross section using decays to $D^*\mu^-X$ final states in pp collisions at sqrt(s) = 7 TeV with the ATLAS detector," Nucl. Phys. B **864**, 341 (2012).
- 59. G. Aad et al. [ATLAS Collaboration], "Search for a standard model Higgs boson in the mass range 200 600-GeV in the $H \to ZZ \to \ell^+\ell^-q\bar{q}$ decay channel with the ATLAS detector," Phys. Lett. B **717**, 70 (2012).
- 60. G. Aad et al. [ATLAS Collaboration], "Measurement of event shapes at large momentum transfer with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C 72, 2211 (2012).
- 61. G. Aad *et al.* [ATLAS Collaboration], "Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 4.7 fb⁻¹ of $\sqrt{s} = 7$ TeV proton-proton collisions," JHEP **1207**, 167 (2012).
- 62. G. Aad et al. [ATLAS Collaboration], "Search for the Standard Model Higgs boson in the $H \to WW(*) \to \ell\nu\ell\nu$ decay mode with 4.7 /fb of ATLAS data at $\sqrt{s} = 7$ TeV," Phys. Lett. B **716**, 62 (2012).
- 63. G. Aad *et al.* [ATLAS Collaboration], "A search for flavour changing neutral currents in top-quark decays in pp collision data collected with the ATLAS detector at $\sqrt{s} = 7$ TeV," JHEP **1209**, 139 (2012).
- 64. G. Aad *et al.* [ATLAS Collaboration], "Search for a Standard Model Higgs boson in the H-i ZZ-i $l^+l^-\nu\bar{\nu}$ decay channel using 4.7 fb^{-1} of sqrt(s) = 7 TeV data with the ATLAS detector," Phys. Lett. B **717**, 29 (2012).
- 65. G. Aad et al. [ATLAS Collaboration], "Evidence for the associated production of a W boson and a top quark in ATLAS at $\sqrt{s} = 7$ TeV," Phys. Lett. B **716**, 142 (2012).
- 66. G. Aad et al. [ATLAS Collaboration], "A search for $t\bar{t}$ resonances with the ATLAS detector in 2.05 fb⁻¹ of proton-proton collisions at $\sqrt{s}=7$ TeV," Eur. Phys. J. C **72**, 2083 (2012).

- 67. ** G. Aad et al. [ATLAS Collaboration], "Measurement of the t-channel single top-quark production cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **717**, 330 (2012).
- 68. G. Aad et al. [ATLAS Collaboration], "Measurement of $W\gamma$ and $Z\gamma$ production cross sections in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous triple gauge couplings with the ATLAS detector," Phys. Lett. B **717**, 49 (2012).
- 69. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the W boson polarization in top quark decays with the ATLAS detector," JHEP **1206**, 088 (2012).
- 70. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the top quark pair cross section with ATLAS in pp collisions at sqrt(s) = 7 TeV using final states with an electron or a muon and a hadronically decaying τ lepton," Phys. Lett. B **717**, 89 (2012).
- 71. G. Aad *et al.* [ATLAS Collaboration], "Search for the resonances in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. Lett. **109**, 081801 (2012).
- 72. G. Aad et al. [ATLAS Collaboration], "Search for lepton flavour violation in the $e\mu$ continuum with the ATLAS detector in $\sqrt{s}=7$ TeV pp collisions at the LHC," Eur. Phys. J. C **72**, 2040 (2012).
- 73. A. Collaboration *et al.* [ATLAS Collaboration], "Search for a fermiophobic Higgs boson in the diphoton decay channel with the ATLAS detector," Eur. Phys. J. C **72**, 2157 (2012).
- 74. G. Aad *et al.* [ATLAS Collaboration], "Search for scalar top quark pair production in natural gauge mediated supersymmetry models with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Phys. Lett. B **715**, 44 (2012).
- 75. G. Aad *et al.* [ATLAS Collaboration], "Measurement of τ polarization in $W > \tau \nu$ decays with the ATLAS detector in pp collisions at sqrt(s) = 7 TeV," Eur. Phys. J. C **72**, 2062 (2012).
- 76. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector," Phys. Rev. Lett. **108**, 261804 (2012).
- 77. G. Aad et al. [ATLAS Collaboration], "Search for TeV-scale gravity signatures in final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 7$ TeV," Phys. Lett. B **716**, 122 (2012).
- 78. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry with jets, missing transverse momentum and at least one hadronically decaying τ lepton in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **714**, 197 (2012).
- 79. G. Aad et al. [ATLAS Collaboration], "Search for charged Higgs bosons decaying via $H^+ \to \tau \nu$ in top quark pair events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1206**, 039 (2012).

- 80. G. Aad et al. [ATLAS Collaboration], "Search for resonant WZ production in the $WZ \to \ell\nu\ell'\ell'$ channel in $\sqrt{s}=7$ TeV pp collisions with the ATLAS detector," Phys. Rev. D 85, 112012 (2012).
- 81. G. Aad *et al.* [ATLAS Collaboration], "Search for pair production of a new quark that decays to a Z boson and a bottom quark with the ATLAS detector," Phys. Rev. Lett. **109**, 071801 (2012).
- 82. G. Aad *et al.* [ATLAS Collaboration], "Search for the decay $B_s^0 \to \mu\mu$ with the ATLAS detector," Phys. Lett. B **713**, 387 (2012).
- 83. G. Aad *et al.* [ATLAS Collaboration], "Search for events with large missing transverse momentum, jets, and at least two τ leptons in 7 TeV proton-proton collision data with the ATLAS detector," Phys. Lett. B **714**, 180 (2012).
- 84. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector and limits on anomalous gauge couplings," Phys. Lett. B **712**, 289 (2012).
- 85. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b^- jets with the ATLAS detector," Phys. Rev. D **85**, 112006 (2012).
- 86. G. Aad *et al.* [ATLAS Collaboration], "Search for gluinos in events with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Phys. Rev. Lett. **108**, 241802 (2012).
- 87. G. Aad et al. [ATLAS Collaboration], "Measurement of the top quark mass with the template method in the $t\bar{t}$ - \bar{t} lepton + jets channel using ATLAS data," Eur. Phys. J. C 72, 2046 (2012).
- 88. G. Aad *et al.* [ATLAS Collaboration], "Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C **72**, 2056 (2012).
- 89. G. Aad et al. [ATLAS Collaboration], "Measurement of $t\bar{t}$ production with a veto on additional central jet activity in pp collisions at sqrt(s) = 7 TeV using the ATLAS detector," Eur. Phys. J. C **72**, 2043 (2012).
- 90. G. Aad *et al.* [ATLAS Collaboration], "Jet mass and substructure of inclusive jets in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS experiment," JHEP **1205**, 128 (2012).
- 91. G. Aad et al. [ATLAS Collaboration], "Measurement of the charge asymmetry in top quark pair production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector," Eur. Phys. J. C **72**, 2039 (2012).
- 92. G. Aad *et al.* [ATLAS Collaboration], "Observation of spin correlation in $t\bar{t}$ events from pp collisions at sqrt(s) = 7 TeV using the ATLAS detector," Phys. Rev. Lett. **108**, 212001 (2012).

- 93. G. Aad et al. [ATLAS Collaboration], "Determination of the strange quark density of the proton from ATLAS measurements of the $W->l\nu$ and Z->ll cross sections," Phys. Rev. Lett. **109**, 012001 (2012).
- 94. G. Aad *et al.* [ATLAS Collaboration], "Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC," JHEP **1205**, 157 (2012).
- 95. G. Aad et al. [ATLAS Collaboration], "Search for second generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C **72**, 2151 (2012).
- 96. G. Aad et al. [ATLAS Collaboration], "Measurement of the production cross section of an isolated photon associated with jets in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 85, 092014 (2012).
- 97. G. Aad *et al.* [ATLAS Collaboration], "Forward-backward correlations and charged-particle azimuthal distributions in pp interactions using the ATLAS detector," JHEP **1207**, 019 (2012).
- 98. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector," Phys. Rev. C **86**, 014907 (2012).
- 99. G. Aad et al. [ATLAS Collaboration], "Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment," Eur. Phys. J. C 72, 2001 (2012).
- 100. G. Aad et al. [ATLAS Collaboration], "Search for a light Higgs boson decaying to long-lived weakly-interacting particles in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. Lett. 108, 251801 (2012).
- 101. G. Aad *et al.* [ATLAS Collaboration], "Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC," Eur. Phys. J. C **73**, 2305 (2013).
- 102. G. Aad et al. [ATLAS Collaboration], "Search for new particles decaying to ZZ using final states with leptons and jets with the ATLAS detector in $\sqrt{s} = 7$ Tev proton-proton collisions," Phys. Lett. B **712**, 331 (2012).
- 103. G. Aad *et al.* [ATLAS Collaboration], "Search for FCNC single top-quark production at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **712**, 351 (2012).
- 104. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the azimuthal ordering of charged hadrons with the ATLAS detector," Phys. Rev. D **86**, 052005 (2012).
- 105. G. Aad *et al.* [ATLAS Collaboration], "Search for down-type fourth generation quarks with the ATLAS detector in events with one lepton and hadronically decaying W bosons," Phys. Rev. Lett. **109**, 032001 (2012).

- 106. G. Aad et al. [ATLAS Collaboration], "Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1204**, 069 (2012).
- 107. G. Aad et al. [ATLAS Collaboration], "Measurement of the cross section for top-quark pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector using final states with two high-pt leptons," JHEP **1205**, 059 (2012).
- 108. G. Aad *et al.* [ATLAS Collaboration], "Search for anomaly-mediated supersymmetry breaking with the ATLAS detector based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C **72**, 1993 (2012).
- 109. G. Aad *et al.* [ATLAS Collaboration], "Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D **86**, 012007 (2012).
- 110. G. Aad *et al.* [ATLAS Collaboration], "Search for pair production of a heavy up-type quark decaying to a W boson and a b quark in the lepton+jets channel with the ATLAS detector," Phys. Rev. Lett. **108**, 261802 (2012).
- 111. G. Aad et al. [ATLAS Collaboration], "Measurement of the inclusive W^{\pm} and Z/gamma cross sections in the electron and muon decay channels in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 85, 072004 (2012).
- 112. G. Aad *et al.* [ATLAS Collaboration], "Search for Diphoton Events with Large Missing Transverse Momentum in 1 fb⁻1 of 7 TeV Proton-Proton Collision Data with the ATLAS Detector," Phys. Lett. B **710**, 519 (2012).
- 113. G. Aad *et al.* [ATLAS Collaboration], "Search for production of resonant states in the photon-jet mass distribution using pp collisions at $\sqrt{s} = 7$ TeV collected by the ATLAS detector," Phys. Rev. Lett. **108**, 211802 (2012).
- 114. G. Aad *et al.* [ATLAS Collaboration], "Jet energy measurement with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C **73**, 2304 (2013).
- 115. G. Aad *et al.* [ATLAS Collaboration], "Measurement of inclusive jet and dijet production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector," Phys. Rev. D **86**, 014022 (2012).
- 116. G. Aad et al. [ATLAS Collaboration], "Search for heavy vector-like quarks coupling to light quarks in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **712**, 22 (2012).
- 117. G. Aad et al. [ATLAS Collaboration], "Observation of a new χ_b state in radiative transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS," Phys. Rev. Lett. 108, 152001 (2012).
- 118. G. Aad *et al.* [ATLAS Collaboration], "Search for contact interactions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **712**, 40 (2012).

- 119. G. Aad *et al.* [ATLAS Collaboration], "Search for scalar bottom pair production with the ATLAS detector in pp Collisions at $\sqrt{s} = 7$ TeV," Phys. Rev. Lett. **108**, 181802 (2012).
- 120. G. Aad *et al.* [ATLAS Collaboration], "Search for Extra Dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector," Phys. Lett. B **710**, 538 (2012).
- 121. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the WZ production cross section and limits on anomalous triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **709**, 341 (2012).
- 122. G. Aad et al. [ATLAS Collaboration], "Search for the Standard Model Higgs boson in the decay channel $H \to ZZ^{(*)} \to 4l$ with 4.8 fb⁻¹ of pp collision data at $\sqrt{s} = 7$ TeV with ATLAS," Phys. Lett. B **710**, 383 (2012).
- 123. G. Aad *et al.* [ATLAS Collaboration], "Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb⁻¹ of pp collisions at \sqrt{s} =7 TeV with ATLAS," Phys. Rev. Lett. **108**, 111803 (2012).
- 124. ** G. Aad *et al.* [ATLAS Collaboration], "Combined search for the Standard Model Higgs boson using up to 4.9 fb⁻¹ of pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC," Phys. Lett. B **710**, 49 (2012).
- 125. G. Aad *et al.* [ATLAS Collaboration], "Rapidity gap cross sections measured with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C **72**, 1926 (2012).
- 126. G. Aad *et al.* [ATLAS Collaboration], "Search for anomalous production of prompt likesign muon pairs and constraints on physics beyond the Standard Model with the ATLAS detector," Phys. Rev. D **88**, 032004 (2012).
- 127. G. Aad *et al.* [ATLAS Collaboration], "Search for first generation scalar leptoquarks in *pp* collisions at sqrts=7 TeV with the ATLAS detector," Phys. Lett. B **709**, 158 (2012).
- 128. G. Aad *et al.* [ATLAS Collaboration], "Measurement of $D^{*\pm}$ meson production in jets from pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector," Phys. Rev. D **85**, 052005 (2012).
- 129. G. Aad *et al.* [ATLAS Collaboration], "Search for the Higgs boson in the $H \to WW^{(*)} \to l\nu l\nu$ decay channel in pp collisions at sqrts = 7 TeV with the ATLAS detector," Phys. Rev. Lett. **108**, 111802 (2012).
- 130. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the production cross section for Z/γ^* in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D **85**, 032009 (2012).
- 131. G. Aad et al. [ATLAS Collaboration], " K_S^0 and Λ production in pp interactions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector at the LHC," Phys. Rev. D 85, 012001 (2012).

- 132. G. Aad *et al.* [ATLAS Collaboration], "Search for strong gravity signatures in same-sign dimuon final states using the ATLAS detector at the LHC," Phys. Lett. B **709**, 322 (2012).
- 133. ** G. Aad *et al.* [ATLAS Collaboration], "A study of the material in the ATLAS inner detector using secondary hadronic interactions," JINST 7, P01013 (2012).
- 134. G. Aad *et al.* [ATLAS Collaboration], "Searches for supersymmetry with the ATLAS detector using final states with two leptons and missing transverse momentum in sqrts = 7 TeV proton-proton collisions," Phys. Lett. B **709**, 137 (2012).
- 135. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the ZZ production cross section and limits on anomalous neutral triple gauge couplings in proton-proton collisions at \sqrt{s} = 7 TeV with the ATLAS detector," Phys. Rev. Lett. **108**, 041804 (2012).
- 136. G. Aad *et al.* [ATLAS Collaboration], "Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data," Eur. Phys. J. C **72**, 1909 (2012).
- 137. G. Aad *et al.* [ATLAS Collaboration], "Search for Massive Colored Scalars in Four-Jet Final States in sqrts=7 TeV proton-proton collisions with the ATLAS Detector," Eur. Phys. J. C **71**, 1828 (2011).
- 138. G. Aad et al. [Atlas Collaboration], "Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using \sqrt{s} =7 TeV pp collisions with the ATLAS detector," JHEP **1111**, 099 (2011).
- 139. G. Aad *et al.* [Atlas Collaboration], "Performance of the ATLAS Trigger System in 2010," Eur. Phys. J. C **72**, 1849 (2012).
- 140. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the inclusive and dijet cross-sections of b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C **71**, 1846 (2011).
- 141. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in sqrts = 7 TeV pp collisions using 1 fb^{-1} of ATLAS data," Phys. Rev. D **85**, 012006 (2012).
- 142. G. Aad et al. [ATLAS Collaboration], "Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions," Phys. Lett. B **710**, 67 (2012).
- 143. G. Aad et al. [ATLAS Collaboration], "Search for the Standard Model Higgs boson in the decay channel $H \to ZZ^{(*)} \to 4l$ with the ATLAS detector," Phys. Lett. B **705**, 435 (2011).
- 144. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the jet fragmentation function and transverse profile in proton-proton collisions at a center-of-mass energy of 7 TeV with the ATLAS detector," Eur. Phys. J. C **71**, 1795 (2011).

- 145. G. Aad *et al.* [ATLAS Collaboration], "Search for New Phenomena in ttbar Events With Large Missing Transverse Momentum in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector," Phys. Rev. Lett. **108**, 041805 (2012).
- 146. G. Aad et al. [ATLAS Collaboration], "Search for the Higgs boson in the $H \to WW \to l\nu jj$ decay channel in pp collisions at sqrts = 7 TeV with the ATLAS detector," Phys. Rev. Lett. 107, 231801 (2011).
- 147. G. Aad et al. [ATLAS Collaboration], "Search for a Standard Model Higgs boson in the $H \to ZZ \to l^+ l^- \nu \bar{\nu}$ decay channel with the ATLAS detector," Phys. Rev. Lett. 107, 221802 (2011).
- 148. G. Aad *et al.* [ATLAS Collaboration], "Search for displaced vertices arising from decays of new heavy particles in 7 TeV *pp* collisions at ATLAS," Phys. Lett. B **707**, 478 (2012).
- 149. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the cross-section for *b*-jets produced in association with a Z boson at \sqrt{s} =7 TeV with the ATLAS detector," Phys. Lett. B **706**, 295 (2012).
- 150. G. Aad et al. [ATLAS Collaboration], "Measurement of the cross section for the production of a W boson in association with b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **707**, 418 (2012).
- 151. G. Aad *et al.* [ATLAS Collaboration], "Measurements of the electron and muon inclusive cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **707**, 438 (2012).
- 152. ** G. Aad *et al.* [ATLAS Collaboration], "Search for New Physics in the Dijet Mass Distribution using 1 fb⁻¹ of pp Collision Data at $\sqrt{s} = 7$ TeV collected by the ATLAS Detector," Phys. Lett. B **708**, 37 (2012).
- 153. G. Aad et al. [ATLAS Collaboration], "Measurement of the Transverse Momentum Distribution of W Bosons in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector," Phys. Rev. D 85, 012005 (2012).
- 154. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the centrality dependence of the charged particle pseudorapidity distribution in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector," Phys. Lett. B **710**, 363 (2012).
- 155. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the pseudorapidity and transverse momentum dependence of the elliptic flow of charged particles in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector," Phys. Lett. B **707**, 330 (2012).
- 156. G. Aad *et al.* [ATLAS Collaboration], "Performance of Missing Transverse Momentum Reconstruction in Proton-Proton Collisions at 7 TeV with ATLAS," Eur. Phys. J. C **72**, 1844 (2012).
- 157. G. Aad et al. [ATLAS Collaboration], "Search for a heavy Standard Model Higgs boson in the channel $H \to ZZ \to llqq$ using the ATLAS detector," Phys. Lett. B **707**, 27 (2012).

- 158. G. Aad *et al.* [ATLAS Collaboration], "A measurement of the ratio of the W and Z cross sections with exactly one associated jet in pp collisions at $\sqrt{s} = 7$ TeV with ATLAS," Phys. Lett. B **708**, 221 (2012).
- 159. G. Aad et al. [ATLAS Collaboration], "Measurement of the $W \to \tau \nu$ Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment," Phys. Lett. B **706**, 276 (2012).
- 160. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s} = 7$ TeV in dilepton final states with ATLAS," Phys. Lett. B **707**, 459 (2012).
- 161. G. Aad et al. [ATLAS Collaboration], "Measurement of the $Z \to \tau\tau$ Cross Section with the ATLAS Detector," Phys. Rev. D 84, 112006 (2011).
- 162. G. Aad et al. [ATLAS Collaboration], "Search for dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. Lett. **107**, 272002 (2011).
- 163. G. Aad et al. [ATLAS Collaboration], "Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector," Phys. Lett. B **705**, 28 (2011).
- 164. G. Aad et al. [ATLAS Collaboration], "Inclusive search for same-sign dilepton signatures in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1110**, 107 (2011).
- 165. G. Aad et al. [ATLAS Collaboration], "Measurement of the inclusive isolated prompt photon cross-section in pp collisions at \sqrt{s} = 7 TeV using 35 pb-1 of ATLAS data," Phys. Lett. B **706**, 150 (2011).
- 166. G. Aad et al. [ATLAS Collaboration], "Search for neutral MSSM Higgs bosons decaying to $\tau^+\tau^-$ pairs in proton-proton collisions at $\sqrt{s}=7$ TeV with the ATLAS detector," Phys. Lett. B **705**, 174 (2011).
- 167. G. Aad et al. [ATLAS Collaboration], "Properties of jets measured from tracks in protonproton collisions at center-of-mass energy $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 84, 054001 (2011).
- 168. ** D. Asner *et al.* [RD42 Collaboration], "Diamond pixel modules," Nucl. Instrum. Meth. A **636**, S125 (2011).
- 169. G. Aad et al. [ATLAS Collaboration], "Measurement of the transverse momentum distribution of Z/γ^* bosons in proton-proton collisions at \sqrt{s} =7 TeV with the ATLAS detector," Phys. Lett. B **705**, 415 (2011).
- 170. G. Aad *et al.* [ATLAS Collaboration], "Measurement of multi-jet cross sections in proton-proton collisions at a 7 TeV center-of-mass energy," Eur. Phys. J. C **71**, 1763 (2011).
- 171. G. Aad et al. [ATLAS Collaboration], "Measurement of dijet production with a veto on additional central jet activity in pp collisions at \sqrt{s} =7 TeV using the ATLAS detector," JHEP **1109**, 053 (2011).

- 172. G. Aad *et al.* [ATLAS Collaboration], "Search for Diphoton Events with Large Missing Transverse Energy with 36 pb⁻¹ of 7 TeV Proton-Proton Collision Data with the ATLAS Detector," Eur. Phys. J. C **71**, 1744 (2011).
- 173. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the isolated di-photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D **85**, 012003 (2012).
- 174. G. Aad et al. [ATLAS Collaboration], "Measurement of the $\Upsilon(1S)$ Production Cross-Section in pp Collisions at $\sqrt{s}=7$ TeV in ATLAS," Phys. Lett. B **705**, 9 (2011).
- 175. G. Aad et al. [ATLAS Collaboration], "Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions," Phys. Lett. B **705**, 294 (2011).
- 176. G. Aad et al. [ATLAS Collaboration], "Search for Heavy Long-Lived Charged Particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV," Phys. Lett. B **703**, 428 (2011).
- 177. G. Aad et al. [ATLAS Collaboration], "Limits on the production of the Standard Model Higgs Boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Eur. Phys. J. C 71, 1728 (2011).
- 178. G. Aad et al. [ATLAS Collaboration], "Measurement of $W\gamma$ and $Z\gamma$ production in proton-proton collisions at \sqrt{s} =7 TeV with the ATLAS Detector," JHEP **1109**, 072 (2011).
- 179. G. Aad et al. [ATLAS Collaboration], "Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with ATLAS," Phys. Rev. Lett. **107**, 041802 (2011).
- 180. G. Aad *et al.* [ATLAS Collaboration], "Search for Contact Interactions in Dimuon Events from pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector," Phys. Rev. D **84**, 011101 (2011).
- 181. G. Aad et al. [ATLAS Collaboration], "Search for pair production of first or second generation leptoquarks in proton-proton collisions at \sqrt{s} =7 TeV using the ATLAS detector at the LHC," Phys. Rev. D 83, 112006 (2011).
- 182. G. Aad et al. [ATLAS Collaboration], "Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/ψ production in proton-proton collisions at $\sqrt{s} = 7$ TeV," Nucl. Phys. B 850, 387 (2011).
- 183. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the Inelastic Proton-Proton Cross-Section at $\sqrt{s}=7$ TeV with the ATLAS Detector," Nature Commun. **2**, 463 (2011).
- 184. G. Aad *et al.* [ATLAS Collaboration], "Search for an excess of events with an identical flavour lepton pair and significant missing transverse momentum in sqrts = 7 TeV proton-proton collisions with the ATLAS detector," Eur. Phys. J. C **71**, 1647 (2011).

- 185. G. Aad et al. [ATLAS Collaboration], "Search for supersymmetric particles in events with lepton pairs and large missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS experiment," Eur. Phys. J. C 71, 1682 (2011).
- 186. G. Aad *et al.* [ATLAS Collaboration], "Search for high mass dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment," Phys. Lett. B **700**, 163 (2011).
- 187. G. Aad et al. [ATLAS Collaboration], "Search for a heavy particle decaying into an electron and a muon with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC," Phys. Rev. Lett. **106**, 251801 (2011).
- 188. G. Aad *et al.* [ATLAS Collaboration], "Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b-jets," Phys. Lett. B **701**, 398 (2011).
- 189. ** G. Aad *et al.* [ATLAS Collaboration], "Search for New Physics in Dijet Mass and Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector," New J. Phys. **13**, 053044 (2011).
- 190. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the Muon Charge Asymmetry from W Bosons Produced in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **701**, 31 (2011).
- 191. G. Aad *et al.* [ATLAS Collaboration], "Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC," Phys. Lett. B **701**, 1 (2011).
- 192. G. Aad *et al.* [ATLAS Collaboration], "Measurements of underlying-event properties using neutral and charged particles in *pp* collisions at 900 GeV and 7 TeV with the ATLAS detector at the LHC," Eur. Phys. J. C **71**, 1636 (2011).
- 193. G. Aad et al. [ATLAS Collaboration], "Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **701**, 50 (2011).
- 194. G. Aad et al. [Atlas Collaboration], "Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions," Phys. Lett. B **701**, 186 (2011).
- 195. ** G. Aad *et al.* [ATLAS Collaboration], "Measurement of Dijet Azimuthal Decorrelations in pp Collisions at \sqrt{s} =7 TeV," Phys. Rev. Lett. **106**, 172002 (2011).
- 196. G. Aad *et al.* [Atlas Collaboration], "Search for supersymmetry using final states with one lepton, jets, and missing transverse momentum with the ATLAS detector in sqrts = 7 TeV *pp* collisons" Phys. Rev. Lett. **106**, 131802 (2011).
- 197. G. Aad *et al.* [ATLAS Collaboration], "Search for Massive Long-lived Highly Ionising Particles with the ATLAS Detector at the LHC," Phys. Lett. B **698**, 353 (2011).

- 198. ** G. Aad *et al.* [ATLAS Collaboration], "Luminosity Determination in pp Collisions at \sqrt{s} =7 TeV Using the ATLAS Detector at the LHC," Eur. Phys. J. C **71**, 1630 (2011).
- 199. G. Aad et al. [Atlas Collaboration], "Study of Jet Shapes in Inclusive Jet Production in pp Collisions at $\sqrt{s} = 7$ TeV using the ATLAS Detector," Phys. Rev. D 83, 052003 (2011).
- 200. G. Aad *et al.* [ATLAS Collaboration], "Measurement of the production cross section for W-bosons in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Lett. B **698**, 325 (2011).
- 201. G. Aad et al. [Atlas Collaboration], "Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," Phys. Rev. D 83, 052005 (2011).
- 202. G. Aad *et al.* [ATLAS Collaboration], "Charged-particle multiplicities in *pp* interactions measured with the ATLAS detector at the LHC," New J. Phys. **13**, 053033 (2011).
- 203. G. Aad *et al.* [ATLAS Collaboration], "Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector," Phys. Rev. Lett. **106**, 121803 (2011).
- 204. G. Aad et al. [Atlas Collaboration], "Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV," Eur. Phys. J. C **71**, 1577 (2011).
- 205. G. Aad et al. [Atlas Collaboration], "Measurement of underlying event characteristics using charged particles in pp collisions at $\sqrt{s} = 900 GeV$ and 7 TeV with the ATLAS detector," Phys. Rev. D 83, 112001 (2011).
- 206. G. Aad *et al.* [The ATLAS Collaboration], "Studies of the performance of the ATLAS detector using cosmic-ray muons," Eur. Phys. J. C **71**, 1593 (2011).
- 207. G. Aad et al. [Atlas Collaboration], "Measurement of the $W \to l\nu$ and $Z/\gamma^* \to ll$ production cross sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector," JHEP **1012**, 060 (2010).
- 208. ** G. Aad *et al.* [ATLAS Collaboration], "Search for Quark Contact Interactions in Dijet Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector," Phys. Lett. B **694**, 327 (2011).
- 209. G. Aad *et al.* [Atlas Collaboration], "Measurement of inclusive jet and dijet cross sections in proton-proton collisions at 7 TeV centre-of-mass energy with the ATLAS detector," Eur. Phys. J. C **71**, 1512 (2011).
- 210. G. Aad *et al.* [ATLAS Collaboration], "Search for New Particles in Two-Jet Final States in 7 TeV Proton-Proton Collisions with the ATLAS Detector at the LHC," Phys. Rev. Lett. **105**, 161801 (2010).

- 211. G. Aad *et al.* [ATLAS Collaboration], "Readiness of the ATLAS Tile Calorimeter for LHC collisions," Eur. Phys. J. C **70**, 1193 (2010).
- 212. G. Aad *et al.* [ATLAS Collaboration], "Commissioning of the ATLAS Muon Spectrometer with Cosmic Rays," Eur. Phys. J. C **70**, 875 (2010).
- 213. ** G. Aad *et al.* [Atlas Collaboration], "Performance of the ATLAS Detector using First Collision Data," JHEP **1009**, 056 (2010).
- 214. G. Aad *et al.* [ATLAS Collaboration], "The ATLAS Simulation Infrastructure," Eur. Phys. J. C **70**, 823 (2010).
- 215. G. Aad *et al.* [ATLAS Collaboration], "The ATLAS Inner Detector commissioning and calibration," Eur. Phys. J. C **70**, 787 (2010).
- 216. G. Aad et al. [ATLAS Collaboration], "Charged-particle multiplicities in pp interactions at $\sqrt{s} = 900$ GeV measured with the ATLAS detector at the LHC," Phys. Lett. B **688**, 21 (2010).
- 217. G. Aad *et al.* [ATLAS Collaboration], "Drift Time Measurement in the ATLAS Liquid Argon Electromagnetic Calorimeter using Cosmic Muons," Eur. Phys. J. C **70**, 755 (2010).
- 218. G. Aad et al. [ATLAS Collaboration], "Measurement of the centrality dependence of J/ψ yields and observation of Z production in lead-lead collisions with the ATLAS detector at the LHC," Phys. Lett. **B697**, 294-312 (2011).
- 219. G. Aad *et al.* [ATLAS Collaboration], "Observation of a Centrality-Dependent Dijet Asymmetry in Lead-Lead Collisions at $\sqrt{S(NN)} = 2.76$ TeV with the ATLAS Detector at the LHC," Phys. Rev. Lett. **105**, 252303 (2010).
- 220. G. Aad et al. [ATLAS Collaboration], "Charged-particle multiplicities in pp interactions at $\sqrt{s} = 900$ GeV measured with the ATLAS detector at the LHC," Phys.Lett. B688 (2010) p21.
- 221. T. Aaltonen *et al.* [CDF Collaboration], "Search for Hadronic Decays of W and Z Bosons in Photon Events in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ -GeV," Phys. Rev. D **80**, 052011 (2009).
- 222. ** V. Cindro et al., "The ATLAS Beam Conditions Monitor," JINST 3, P02004 (2008).
- 223. ** G. Aad *et al.* [ATLAS Collaboration], "The ATLAS Experiment at the CERN Large Hadron Collider," JINST **3**, S08003 (2008).
- 224. ** M. Mathes, M. Cristinziani, H. Kagan, S. Smith, W. Trischuk, J. Velthuis and N. Wermes, "Characterization of a Single Crystal Diamond Pixel Detector in a High Energy Particle Beam," JINST 3, P12002 (2008).
- 225. G. Aad *et al.* [ATLAS Collaboration], "Expected Performance of the ATLAS Experiment Detector, Trigger and Physics," arXiv:0901.0512.

- 226. J. J. Velthuis *et al.*, "Radiation hard diamond pixel detectors," Nucl. Instrum. Meth. A **591**, 221 (2008).
- 227. ** T. Aaltonen *et al.* [CDF-II Collaboration], "First Run II Measurement of the W Boson Mass," Phys. Rev. D **77**, 112001 (2008).
- 228. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Maximal Flavor Violating Scalars in Same-Charge Lepton Pairs in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV," Phys. Rev. Lett. **102**, 041801 (2009).
- 229. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Third Generation Vector Leptoquarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **77**, 091105 (2008).
- 230. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for resonant $t\bar{t}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **100**, 231801 (2008).
- 231. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for chargino-neutralino production in $p \bar{p}$ collisions at 1.96 TeV with high p_T leptons," Phys. Rev. D **77**, 052002 (2008).
- 232. T. Aaltonen *et al.* [CDF-II Collaboration], "Observation of Exclusive Dijet Production at the Fermilab Tevatron $p \bar{p}$ Collider," Phys. Rev. D **77**, 052004 (2008).
- 233. T. Aaltonen *et al.* [CDF-II Collaboration], "First Flavor-Tagged Determination of Bounds on Mixing-Induced CP Violation in $B_s \to J/\psi \phi$ Decays," Phys. Rev. Lett. **100**, 161802 (2008).
- 234. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for New Heavy Particles Decaying to $Z^0Z^0 \rightarrow eeee$ in p \bar{p} Collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **78**, 012008 (2008).
- 235. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Heavy Top-like Quarks $t' \to Wq$ Using Lepton Plus Jets Events in 1.96 TeV Proton-Antiproton Collisions," Phys. Rev. Lett. **100**, 161803 (2008).
- 236. T. Aaltonen *et al.* [CDF-II Collaboration], "Measurement of Ratios of Fragmentation Fractions for Bottom Hadrons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D 77, 072003 (2008).
- 237. T. Aaltonen *et al.* [CDF-II Collaboration], "First Measurement of ZZ Production in panti-p Collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **100**, 201801 (2008).
- 238. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for the Higgs boson in events with missing transverse energy and b quark jets produced in proton-antiproton collisions at \sqrt{s} =1.96 TeV," Phys. Rev. Lett. **100**, 211801 (2008).
- 239. T. Aaltonen *et al.* [CDF-II Collaboration], "Two-Particle Momentum Correlations in Jets Produced in ppbar Collisions at $\sqrt{s} = 1.96$ -TeV,", Phys. Rev. D **77**, 092001 (2008)
- 240. L. Brigliadori et al. [CDF-II Collaboration], "Search for Pair Production of Scalar Top Quarks Decaying to a τ Lepton and a b Quark in $p \bar{p}$ Collisions at sqrts=1.96 TeV," Phys. Rev. Lett. **101**, 071802 (2008).

- 241. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Heavy, Long-Lived Neutralinos that Decay to Photons at CDF II Using Photon Timing, Phys. Rev. D **78**, 032015 (2008).
- 242. J. C. Armitage *et al.*, "Electron signals in the forward calorimeter prototype for ATLAS," JINST **2**, P11001 (2007).
- 243. T. Aaltonen *et al.* [CDF-II Collaboration], "Measurement of the cross section for W-boson production in association with jets in ppbar collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. D 77, 011108 (2008).
- 244. T. Aaltonen *et al.* [CDF-II Collaboration], "Measurement of Inclusive Jet Cross Sections in $Z/g * *(\rightarrow ee) + \text{ jets Production in } p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **100**, 102001 (2008).
- 245. ** M. Mikuz *et al.*, "Diamond pad detector telescope for beam conditions and luminosity monitoring in ATLAS," Nucl. Instrum. Meth. A **579**, 788 (2007).
- 246. T. Aaltonen *et al.* [CDF-II Collaboration], "First Measurement of the Production of a W Boson in Association with a Single Charm Quark in Proton Anti-proton Collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **100**, 091803 (2008).
- 247. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Standard Model Higgs Bosons Produced in Association with W Bosons," Phys. Rev. Lett. **100**, 041801 (2008).
- 248. T. Aaltonen *et al.* [The CDF Collaboration], "Search for Doubly Charged Higgs Bosons with Lepton-Flavor-Violating Decays involving Tau Leptons," Phys. Rev. Lett. **101**, 121801 (2008).
- 249. T. Aaltonen *et al.* [CDF-II Collaboration], "Observation of Orbitally Excited B_s Mesons," Phys. Rev. Lett. **100**, 082001 (2008).
- 250. T. Aaltonen et al. [CDF-II Collaboration], "A direct Measurement of the W Boson Width in $p\bar{p}$ Collisions at $\sqrt{(s=1.96~\text{TeV},"\text{Phys. Rev. Lett. }\textbf{100},~071801~(2008)}$.
- 251. T. Aaltonen *et al.* [CDF-II Collaboration], "Cross Section Constrained Top Quark Mass Measurement from Dilepton Events at the Tevatron," Phys. Rev. Lett. **100**, 062005 (2008).
- 252. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for Direct Pair Production of Supersymmetric Top and Supersymmetric Bottom Quarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **76**, 072010 (2007).
- 253. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for new physics in high mass electron-positron events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **99**, 171802 (2007).
- 254. T. Aaltonen et al. [CDF-II Collaboration], "Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions," Phys. Rev. Lett. **99**, 242002 (2007).

- 255. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for chargino-neutralino production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **99**, 191806 (2007).
- 256. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for a high-mass diphoton state and limits on Randall-Sundrum gravitons at CDF," Phys. Rev. Lett. **99**, 171801 (2007).
- 257. T. Aaltonen *et al.* [CDF-II Collaboration], "Observation and mass measurement of the baryon Xi_b^- ," Phys. Rev. Lett. **99**, 052002 (2007).
- 258. ** A. Gorisek *et al.*, "ATLAS diamond beam condition monitor," Nucl. Instrum. Meth. A **572**, 67 (2007).
- 259. ** T. Aaltonen *et al.* [CDF-II Collaboration], "First Measurement of the W Boson Mass in Run II of the Tevatron," Phys. Rev. Lett. **99**, 151801 (2007).
- 260. T. Aaltonen *et al.* [CDF-II Collaboration], "First observation of heavy baryons Σ_b and Σ_b^* ," Phys. Rev. Lett. **99**, 202001 (2007).
- 261. T. Aaltonen *et al.* [CDF-II Collaboration], "Measurement of the $p\bar{p} \to t\bar{t}$ production cross- section and the top quark mass at $\sqrt{s} = 1.96$ -TeV in the all-hadronic decay mode," Phys. Rev. D **76**, 072009 (2007).
- 262. T. Aaltonen *et al.* [CDF-II Collaboration], "Search for New Particles Leading to Z+ jets Final States in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **76**, 072006 (2007).
- 263. T. Aaltonen *et al.* [CDF-II Collaboration], "Limits on Anomalous Triple Gauge Couplings in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **76**, 111103 (2007).
- 264. T. Aaltonen *et al.* [CDF-II Collaboration], "Measurement of the top-quark mass using missing E_T + jets events with secondary vertex b-tagging at CDF II," Phys. Rev. D **75**, 111103 (2007).
- 265. A. Abulencia *et al.* [CDF-II Collaboration], "Search for heavy, long-lived particles that decay to photons at CDF II," Phys. Rev. Lett. **99**, 121801 (2007).
- 266. A. Abulencia et al. [CDF-II Collaboration], "Polarization of J/ψ and ψ_{2S} mesons produced in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **99**, 132001 (2007).
- 267. A. Abulencia *et al.* [CDF-II Collaboration], "Precise measurement of the top quark mass in the lepton+jets topology at CDF II," Phys. Rev. Lett. **99**, 182002 (2007).
- 268. A. Abulencia et al. [CDF-II Collaboration], "Measurement of $\sigma_{\chi_{c2}}\mathcal{B}(\chi_{c2} \to J/\psi\gamma)/\sigma_{\chi_{c1}}\mathcal{B}(\chi_{c1} \to J/\psi\gamma)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **98**, 232001 (2007).
- 269. A. Abulencia *et al.* [CDF-II Collaboration], "First Measurement of the Ratio of Central-Electron to Forward-Electron W Partial Cross Sections in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV," Phys. Rev. Lett. **98**, 251801 (2007).
- 270. A. Abulencia *et al.* [CDF-II Collaboration], "Observation of WZ Production," Phys. Rev. Lett. **98**, 161801 (2007).

- 271. A. Abulencia *et al.* [CDF-II Collaboration], "Analysis of the quantum numbers J(PC) of the X(3872)," Phys. Rev. Lett. **98**, 132002 (2007).
- 272. A. Abulencia *et al.* [CDF-II Collaboration], "Measurements of inclusive W and Z cross sections in *pbarp* collisions at $\sqrt{s} = 1.96$ TeV," J. Phys. G **34**, 2457 (2007).
- 273. A. Abulencia *et al.* [CDF-II Collaboration], "Inclusive search for new physics with like-sign dilepton events in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **98**, 221803 (2007).
- 274. A. Abulencia *et al.* [CDF-II Collaboration], "Search for new physics in lepton + photon + X events with 929 pb⁻¹ of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **75**, 112001 (2007).
- 275. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the Inclusive Jet Cross Section using the k_T algorithm in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV with the CDF II Detector," Phys. Rev. D **75**, 092006 (2007), [Erratum-ibid. D **75**, 119901 (2007)].
- 276. A. Abulencia et al. [CDF-II Collaboration], "Search for anomalous production of multilepton events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **98**, 131804 (2007).
- 277. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the Top Quark Mass in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV using the Decay Length Technique," Phys. Rev. D 75, 071102 (2007).
- 278. A. Abulencia *et al.* [CDF-II Collaboration], "Precision measurement of the top quark mass from dilepton events at CDF II,", Phys. Rev. D **75**, 031105 (2007).
- 279. T. Aaltonen *et al.*, [CDF-II Collaboration] "Measurement of the top-quark mass in all-hadronic decays in $p\bar{p}$ collisions at CDF II," Phys. Rev. Lett. **98**, 142001 (2007).
- 280. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the helicity fractions of W bosons from top quark decays using fully reconstructed $t\bar{t}$ events with CDF II," Phys. Rev. D **75**, 052001 (2007).
- 281. A. Abulencia et al. [CDF-II Collaboration], "Search for W' boson decaying to electron-neutrino pairs in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **75**, 091101 (2007).
- 282. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the Ratios of Branching Fractions $\mathcal{B}(\mathcal{B}_{f} \to \mathcal{D}_{f}\pi\pi\pi)/\mathcal{B}(\mathcal{B}' \to \mathcal{D}'\pi\pi\pi)$ and $\mathcal{B}(\mathcal{B}_{f} \to \mathcal{D}_{f}\pi)/\mathcal{B}(\mathcal{B}' \to \mathcal{D}'\pi)$," Phys. Rev. Lett. **98**, 061802 (2007).
- 283. A. Abulencia *et al.* [CDF-II Collaboration], "Observation of $B^0_s \to \psi(2S)\phi$ and measurement of ratio of branching fractions $\mathcal{B}(B^0_s \to \psi(2S)\phi)/\mathcal{B}(B^0_s \to J/\psi\phi)$," Phys. Rev. Lett. **96**, 231801 (2006).
- 284. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the Λ_b^0 lifetime in $\Lambda_b^0 \to J/\psi \Lambda^0$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **98**, 122001 (2007).

- 285. A. Abulencia et al. [CDF-II Collaboration], "Search for V + A current in top quark decay in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **98**, 072001 (2007).
- 286. A. Abulencia *et al.* [CDF-II Collaboration], "Search for Exotic S=-2 Baryons in proton-antiproton Collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. D **75**, 032003 (2007).
- 287. A. Abulencia *et al.* [CDF-II Collaboration], "Observation of Exclusive Electron-Positron Production in Hadron-Hadron Collisions," Phys. Rev. Lett. **98**, 112001 (2007).
- 288. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the B^+ production cross-section in $p\bar{p}$ collisions at $\sqrt{s}=1960\text{-GeV}$," Phys. Rev. D **75**, 012010 (2007).
- 289. ** W. Adam *et al.* [RD42 Collaboration], "Radiation hard diamond sensors for future tracking applications," Nucl. Instrum. Meth. A **565**, 278 (2006).
- 290. ** A. Abulencia *et al.* [CDF-II Collaboration], "Observation of B_s^0 \bar{B}_s^0 Oscillations," Phys. Rev. Lett. **97**, 242003 (2006).
- 291. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV in the All Hadronic Decay Mode," Phys. Rev. D **74**, 072005 (2006).
- 292. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV using Lepton + Jets Events with Jet Probability b tagging," Phys. Rev. D **74**, 072006 (2006).
- 293. A. Abulencia *et al.* [CDF-II Collaboration], "Observation of $B_s^0 \to K^+K^-$ and Measurements of Branching Fractions of Charmless Two-body Decays of B^0 and B_s^0 Mesons in $\bar{p}p$ Collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **97**, 211802 (2006).
- 294. A. Abulencia et al. [CDF-II Collaboration], "Search for excited and exotic muons in the $\mu\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. 97, 191802 (2006).
- 295. ** A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the B_s^0 \bar{B}_s^0 Oscillation Frequency," Phys. Rev. Lett. **97**, 062003 (2006).
- 296. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **97**, 082004 (2006).
- 297. A. Abulencia et al. [CDF-II Collaboration], "Search for a neutral Higgs boson decaying to a W boson pair in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. 97, 081802 (2006).
- 298. A. Abulencia *et al.* [CDF-II Collaboration], "Top quark mass measurement from dilepton events at CDF-II with the matrix-element method," Phys. Rev. D **74**, 032009 (2006).
- 299. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the b jet cross-section in events with a Z boson in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **74**, 032008 (2006).

- 300. A. Abulencia et al. [CDF-II Collaboration], "Search for new physics in lepton + photon + X events with 305 pb⁻¹ of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **97**, 031801 (2006).
- 301. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the ratio of branching fractions $\mathcal{B}(\mathcal{D}' \to \mathcal{K}^+\pi^-)/\mathcal{B}(\mathcal{D}' \to \mathcal{K}^-\pi^+)$ using the CDF II Detector," Phys. Rev. D **74**, 031109 (2006).
- 302. Z. Natkaniec *et al.*, [BELLE SVD Group], "Status of the Belle silicon vertex detector," Nucl. Instrum. Meth. A **560**, 1 (2006).
- 303. A. Abulencia *et al.* [CDF-II Collaboration], "Search for large extra dimensions in the production of jets and missing transverse energy in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **97**, 171802 (2006).
- 304. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV using missing E(t) + jets events with secondary vertex b-tagging," Phys. Rev. Lett. **96**, 202002 (2006).
- 305. A. Abulencia et al. [CDF-II Collaboration], "Measurement of the B_c^+ meson lifetime using $B_c^+ \to J/\psi e^+ \nu/e$," Phys. Rev. Lett. 97, 012002 (2006).
- 306. A. Abulencia et al. [CDF-II Collaboration], "Search for high-mass resonances decaying to $e \mu$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **96**, 211802 (2006).
- 307. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the top quark mass using template methods on dilepton events in proton antiproton collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **73**, 112006 (2006).
- 308. ** H. Aihara *et al.*, [BELLE SVD Group], "Belle SVD2 vertex detector," Nucl. Instrum. Meth. A **568**, 269 (2006).
- 309. A. Abulencia et al. [CDF-II Collaboration], "Search for $Z' \to e^+e^-$ using dielectron mass and angular distribution," Phys. Rev. Lett. **96**, 211801 (2006).
- 310. A. Abulencia et al. [CDF-II Collaboration], "Search for $H \to b\bar{b}$ produced in association with W bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **96** (2006) 081803.
- 311. A. Abulencia et al. [CDF-II Collaboration], "Search for charged Higgs bosons from top quark decays in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV," Phys. Rev. Lett. **96** (2006) 042003.
- 312. A. Abulencia *et al.* [CDF-II Collaboration], "Precision top quark mass measurement in the lepton + jets topology in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **96** (2006) 022004.
- 313. A. Abulencia *et al.* [CDF-II Collaboration], "Search for neutral MSSM Higgs bosons decaying to τ pairs in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV," Phys. Rev. Lett. **96** (2006) 011802.

- 314. A. Abulencia *et al.* [CDF-II Collaboration], "Search for $B_s^0 \to \mu^+\mu^-$ and $B_d^0 \to \mu^+\mu^-$ decays in $p\bar{p}$ collisions with CDF II," Phys. Rev. Lett. **95** (2005) 221805, [Erratum-ibid. **95** (2005) 249905.
- 315. A. Abulencia *et al.* [CDF-II Collaboration], "Search for new high mass particles decaying to lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **95** (2005) 252001
- 316. D. Acosta *et al.* [CDF-II Collaboration], "Search for $\Lambda_b \to p\pi and\Lambda_b \to pK$ decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. D **72** (2005) 051104.
- 317. D. Acosta *et al.* [CDF-II Collaboration], "Search for W and Z bosons in the reaction $\bar{p}p \to 2$ jets + γ at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **73** (2006) 012001.
- 318. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the dipion mass spectrum in $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decays," Phys. Rev. Lett. **96**, 102002 (2006).
- 319. A. Abulencia *et al.* [CDF-II Collaboration], "A search for scalar bottom quarks from gluino decays in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **96**, 171802 (2006).
- 320. A. Abulencia *et al.* [CDF-II Collaboration], "Top quark mass measurement from dilepton events at CDF II," Phys. Rev. Lett. **96**, 152002 (2006).
- 321. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of mass and width of the excited charmed meson states D_1^0 and D_2^{*0} ," Phys. Rev. D **73**, 051104 (2006).
- 322. A. Abulencia *et al.* [CDF-II Collaboration], "Search for anomalous semileptonic decay of heavy flavor hadrons produced in association with a W boson at CDF II," Phys. Rev. D **73**, 051101 (2006).
- 323. A. Abulencia *et al.* [CDF-II II Collaboration], "Measurement of the inclusive jet cross section using the k_t algorithm in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **96**, 122001 (2006).
- 324. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the inclusive jet cross section in $p\bar{p}$ interactions at $\sqrt{s}=1.96$ TeV using a cone-based jet algorithm," Phys. Rev. D **74**, 071103 (2006).
- 325. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the top quark mass with the dynamical likelihood method using lepton plus jets events with *b*-tags in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. D **73**, 092002 (2006).
- 326. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the helicity of W bosons in top-quark decays," Phys. Rev. D **73**, 111103 (2006).
- 327. A. Abulencia et al. [CDF-II Collaboration], "A search for $t \to \tau \nu q$ in $t\bar{t}$ production," Phys. Lett. B **639**, 172 (2006).
- 328. A. Abulencia *et al.* [CDF-II Collaboration], "Top quark mass measurement using the template method in the lepton + jets channel at CDF II," Phys. Rev. D **73**, 032003 (2006).

- 329. A. Abulencia *et al.* [CDF-II Collaboration], "Direct search for Dirac magnetic monopoles in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. Lett. **96**, 201801 (2006).
- 330. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of *b* hadron masses in exclusive J/ψ decays with the CDF detector," Phys. Rev. Lett. **96**, 202001 (2006).
- 331. A. Abulencia *et al.* [CDF-II Collaboration], "Measurement of the ratios of branching fractions $\mathcal{B}(B_s^0 \to D_s^- \pi^+)/\mathcal{B}(B^0 \to D^- \pi^+)$ and $\mathcal{B}(B^+ \to \bar{D}^0 \pi^+)/\mathcal{B}(B^0 \to D^- \pi^+)$," Phys. Rev. Lett. **96**, 191801 (2006).
- 332. A. Abulencia *et al.* [CDF-II Collaboration], "Search for second-generation scalar leptoquarks in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **73**, 051102 (2006).
- 333. D. Acosta *et al.* [CDF-II Collaboration], "Search for first-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV," Phys. Rev. D **72** (2005) 051107.
- 334. D. Acosta *et al.* [CDF-II Collaboration], "A search for supersymmetric Higgs bosons in the di- τ decay mode in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **72** (2005) 072004.
- 335. D. Acosta *et al.* [CDF-II Collaboration], "Search for new physics using high mass τ pairs from 1.96 TeV $p\bar{p}$ collisions," Phys. Rev. Lett. **95** (2005) 131801.
- 336. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV using lepton plus jets events with semileptonic B decays to muons," Phys. Rev. D **72** (2005) 032002.
- 337. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of $\mathcal{B}(t \to Wb)/\mathcal{B}(t \to Wq)$ at the Collider Detector at Fermilab," Phys. Rev. Lett. **95** (2005) 102002.
- 338. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the cross section for $t\bar{t}$ production in $p\bar{p}$ collisions using the kinematics of lepton + jets events," Phys. Rev. D **72** (2005) 052003.
- 339. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **73** (2006) 052002.
- 340. D. Acosta *et al.* [CDF-II Collaboration], "Search for Higgs bosons decaying into $b\bar{b}$ and produced in association with a vector boson in proton antiproton collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. Lett. **95** (2005) 051801.
- 341. D. Acosta *et al.* [CDF-II Collaboration], "Search for Z Z and Z W production in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **71** (2005) 091105.
- 342. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the W^+W^- production cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV using dilepton events," Phys. Rev. Lett. **94** (2005) 211801.
- 343. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the moments of the hadronic invariant mass distribution in semileptonic *B* decays," Phys. Rev. D **71** (2005) 051103.

- 344. D. Acosta *et al.* [CDF-II Collaboration], "First evidence for $B_s^0 \to \phi \phi$ decay and measurements of branching ratio and A(CP) for $B^+ \to \phi K^+$," Phys. Rev. Lett. **95** (2005) 031801.
- 345. D. Acosta *et al.* [CDF-II Collaboration], "Search for long-lived doubly-charged Higgs bosons in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **95** (2005) 071801.
- 346. T. Tsuboyama *et al.*, [BELLE SVD Group], "The Silicon Vertex Detector for the super B factory," Nucl. Instrum. Meth. A **541** (2005) 421.
- 347. R. Stamen *et al.* [BELLE SVD Group], "Status of the Belle Silicon Vertex Detector," Nucl. Instrum. Meth. A **541** (2005) 61.
- 348. R. Abe *et al.* [BELLE SVD Group], "The new beampipe for the Belle experiment," Nucl. Instrum. Meth. A **535**, 558 (2004).
- 349. R. Abe *et al.* [BELLE SVD Group], "BELLE/SVD2 status and performance," Nucl. Instrum. Meth. A **535**, 379 (2004).
- 350. H. Ishino *et al.* [BELLE SVD Group], "The data acquisition system of the Belle silicon vertex detector (SVD) upgrade," IEEE Trans. Nucl. Sci. **51**, 2064 (2004).
- 351. ** T. Ziegler *et al.* [BELLE SVD Group], "The Belle trigger system with the new silicon vertex detector SVD2," IEEE Trans. Nucl. Sci. **51**, 1852 (2004).
- 352. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the forward-backward charge asymmetry from $W \to e\nu$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **71**, 051104 (2005).
- 353. ** W. Adam *et al.* [RD42 Collaboration], "New developments in CVD diamond for detector applications," Eur. Phys. J. C **33**, S1014 (2004).
- 354. ** D. Acosta *et al.* [CDF-II Collaboration], "Search for electroweak single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **71**, 012005 (2005).
- 355. D. Acosta *et al.* [CDF-II Collaboration], "Search for excited and exotic electrons in the $e\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **94**, 101802 (2005).
- 356. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of $W\gamma$ and $Z\gamma$ production in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **94**, 041803 (2005).
- 357. M. Friedl *et al.* [BELLE SVD Group], "Readout, First- and Second-Level Triggers of the new Belle Silicon Vertex Detector," Nucl. Instrum. Meth. A **535**, 491 (2004).
- 358. ** T. Lari, A. Oh, N. Wermes, H. Kagan, M. Keil and W. Trischuk, "Characterization and Modeling of Non-Uniform Charge Collection in CVD Diamond Pixel Detectors," Nucl. Instrum. Meth. A **537**, 581 (2005).
- 359. D. Acosta *et al.* [CDF-II Collaboration], "First measurements of inclusive W and Z cross sections from Run II of the Tevatron collider," Phys. Rev. Lett. **94**, 091803 (2005).

- 360. D. Acosta *et al.* [CDF Collaboration], "Comparison of three-jet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV to predictions from a next-to-leading order QCD calculation," Phys. Rev. D **71**, 032002 (2005).
- 361. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of partial widths and search for direct CP violation in D^0 meson decays to K^-K^+ and $\pi^-\pi^+$," Phys. Rev. Lett. **94**, 122001 (2005).
- 362. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV using kinematic fitting of b-tagged lepton + jet events," Phys. Rev. D **71**, 072005 (2005).
- 363. D. Acosta *et al.* [CDF-II Collaboration], "Search for doubly-charged Higgs bosons decaying to dileptons in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **93**, 221802 (2004).
- 364. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the J/ψ meson and *b*-hadron production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1960$ -GeV," Phys. Rev. D **71**, 032001 (2005).
- 365. D. Acosta *et al.* [CDF-II Collaboration], "Analysis of decay-time dependence of angular distributions in $B_s^0 \to J/\psi \phi$ and $B_d^0 \to J/\psi K^{*0}$ decays and measurement of the lifetime difference between B_s mass eigenstates," Phys. Rev. Lett. **94**, 101803 (2005).
- 366. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the cross section for prompt diphoton production in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **95**, 022003 (2005).
- 367. D. Acosta *et al.* [CDF-II Collaboration], "Search for anomalous kinematics in $t\bar{t}$ dilepton events at CDF II," Phys. Rev. Lett. **95**, 022001 (2005).
- 368. D. Acosta *et al.* [CDF Collaboration], "Measurements of bottom anti-bottom azimuthal production correlations in proton antiproton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **71**, 092001 (2005).
- 369. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the forward-backward charge asymmetry of electron positron pairs in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," arXiv:hep-ex/0411059.
- 370. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of charged particle multiplicities in gluon and quark jets in proton anti-proton collisions at $\sqrt{s} = 1.8$ -TeV," FERMILAB-PUB-04-113-E.
- 371. D. Acosta *et al.* [CDF-II Collaboration], "Search for scalar leptoquark pairs decaying to $\nu\bar{\nu}q\bar{q}$ in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. D **71**, 112001 (2005), [Erratum-ibid. D **71**, 119901 (2005)].

- 372. D. Acosta *et al.* [CDF-II Collaboration], "Search for anomalous production of diphoton events with missing transverse energy at CDF and limits on gauge-mediated supersymmetry-breaking models," Phys. Rev. D **71**, 031104 (2005).
- 373. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV using lepton + jets events with secondary vertex *b*-tagging," Phys. Rev. D **71**, 052003 (2005).
- 374. ** W. Adam *et al.* [RD42 Collaboration], "The development of diamond tracking detectors for the LHC," Nucl. Instrum. Meth. A **514**, 79 (2003).
- 375. D. Acosta *et al.* [CDF-II Collaboration], "Inclusive search for anomalous production of high p_T like-sign lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **93**, 061802 (2004).
- 376. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV using dilepton events," Phys. Rev. Lett. **93**, 142001 (2004).
- 377. D. Acosta *et al.* [CDF-II Collaboration], "Direct photon cross section with conversions at CDF," Phys. Rev. D **70**, 074008 (2004).
- 378. V. M. Abazov *et al.* [CDF Collaboration], "Combination of CDF and D0 results on W boson mass and width," Phys. Rev. D **70**, 092008 (2004).
- 379. D. Acosta *et al.* [CDF Collaboration], "Measurement of the W boson polarization in top decay at CDF at \sqrt{s} = 1.8-TeV," Phys. Rev. D **71**, 031101 (2005) [Erratum-ibid. D **71**, 059901 (2005)].
- 380. D. Acosta *et al.* [CDF-II Collaboration], "The underlying event in hard interactions at the Tevatron anti-p p collider," Phys. Rev. D **70**, 072002 (2004).
- 381. D. Acosta *et al.* [CDF-II Collaboration], "Search for $B^0_s \to \mu^+\mu^-$ and $B^0_d \to \mu^+\mu^-$ decays in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ -TeV," Phys. Rev. Lett. **93**, 032001 (2004).
- 382. D. Acosta *et al.* [CDF-II Collaboration], "Observation of the narrow state X(3872) \rightarrow J/psi pi+ pi- in anti-p p collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **93**, 072001 (2004).
- 383. D. Acosta *et al.* [CDF Collaboration], "Measurement of the polar-angle distribution of leptons from W boson decay as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **70**, 032004 (2004).
- 384. D. Acosta *et al.* [CDF Collaboration], "Inclusive double pomeron exchange at the Fermilab Tevatron $\bar{p}p$ collider," Phys. Rev. Letter **93**, 141601 (2004).
- 385. ** D. Acosta *et al.* [CDF-II Collaboration], "Optimized search for single top quark production at the Fermilab Tevatron," Phys. Rev. D **69**, 052003 (2004).
- 386. ** W. Adam *et al.* [RD42 Collaboration], "Status Of The R & D Activity On Diamond Particle Detectors," Nucl. Instrum. Meth. A **511**, 124 (2003).

- 387. D. Acosta *et al.* [CDF Collaboration], "Measurement of the average time-integrated mixing probability of *b*-flavored hadrons produced at the Tevatron," Phys. Rev. D **69**, 012002 (2004)
- 388. D. Acosta *et al.* [CDF-II Collaboration], "Search for the flavour-changing neutral current decay $D^0 \to \mu + \mu \text{in } p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. D **68**, 091101 (2003).
- 389. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of the mass difference $m(D_s^+) m(D^+)$ at CDF II," Phys. Rev. D **68**, 072004 (2003).
- 390. D. Acosta *et al.* [CDF-II Collaboration], "Measurement of prompt charm meson production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV," Phys. Rev. Lett. **91**, 241804 (2003)
- 391. D. Acosta *et al.* [CDF Collaboration], "Search for pair production of scalar top quarks in R-parity violating decay modes in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **92**, 051803 (2004).
- 392. D. Acosta *et al.* [CDF Collaboration], "Search for associated production of Υ and vector boson in $p\bar{p}$ collisions At $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **90**, 221803 (2003).
- 393. ** M. Keil *et al.* [RD42 Collaboration], "New results on Diamond pixel sensors using ATLAS front end electronics," Nucl. Instrum. Meth. A **501**, 153 (2003).
- 394. D. Acosta *et al.* [CDF Collaboration], "Search for the supersymmetric partner of the top quark in dilepton events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **90**, 251801 (2003).
- 395. ** W. Trischuk [BELLE Collaboration], "Measurement of CP asymmetries at BELLE," eConf **C020805**, TW02 (2002) [arXiv:hep-ex/0212059].
- 396. D. Acosta *et al.* [CDF Collaboration], "Cross section for forward J/ψ production in $p\bar{p}$ collisions at \sqrt{s} =1.8 TeV," Phys. Rev. D **66**, 092001 (2002).
- 397. D. Acosta *et al.* [CDF Collaboration], "Search for long-lived charged massive particles in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **90**, 131801 (2003).
- 398. D. Acosta *et al.* [CDF Collaboration], "Search for a W' boson decaying to a top and bottom quark pair in 1.8-TeV $p\bar{p}$ collisions," Phys. Rev. Lett. **90**, 081802 (2003).
- 399. D. Acosta *et al.* [CDF Collaboration], "Search for radiative *b*-hadron decays in $p\bar{p}$ collisions at $\sqrt{s}=1.8$ -TeV," Phys. Rev. D **66**, 112002 (2002).
- 400. ** J. Kaneko *et al.* [BELLE Collaboration], "Measurement of the electroweak penguin process $B \to X_s l^+ l^-$," Phys. Rev. Lett. **90**, 021801 (2003).
- 401. D. Acosta *et al.* [CDF Collaboration], "Momentum distribution of charged particles in jets in dijet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV and comparisons to perturbative QCD predictions," Phys. Rev. D **68**, 012003 (2003).

- 402. ** K. Abe *et al.* [BELLE Collaboration], "An improved measurement of mixing-induced CP violation in the neutral *B* meson system," Phys. Rev. D **66**, 071102 (2002)
- 403. D. Acosta *et al.* [CDF Collaboration], "Limits on extra dimensions and new particle production in the exclusive photon and missing energy signature in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **89**, 281801 (2002).
- 404. T. Affolder *et al.* [CDF Collaboration], "Charged jet evolution and the underlying event in proton-antiproton collision \sqrt{s} at 1.8 TeV," Phys. Rev. D **65**, 092002 (2002).
- 405. D. Acosta *et al.* [CDF Collaboration], section $d\sigma/dp_T$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **65**, 052005 (2002).
- 406. ** W. Adam *et al.* [RD42 Collaboration], "Radiation tolerance of CVD diamond detectors for pions and protons," Nucl. Instrum. Meth. A **476**, 686 (2002).
- 407. D. Acosta *et al.* [CDF Collaboration], "Measurement of *B*-meson lifetimes using fully reconstructed *B* decays produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **65**, 092009 (2002).
- 408. ** K. Abe *et al.* [BELLE Collaboration], "Observation of double $c\bar{c}$ production in e^+e^- annihilation at \sqrt{s} 10.6-GeV," Phys. Rev. Lett. **89**, 142001 (2002).
- 409. D. Acosta *et al.* [CDF Collaboration], "Branching ratio measurements of exclusive B^+ decays to charmonium with the Collider Detector at Fermilab," Phys. Rev. D **66**, 052005 (2002).
- 410. D. Acosta *et al.* [CDF Collaboration], "Measurement of the ratio of *b* quark production cross sections in $\bar{p}p$ collisions at $\sqrt{s} = 630$ -GeV and $\sqrt{s} = 1800$ -GeV," Phys. Rev. D **66**, 032002 (2002).
- 411. ** K. F. Chen *et al.* [BELLE Collaboration], "Measurement of CP-violating parameters in $B \to \eta' K$ decays," Phys. Lett. B **546**, 196 (2002).
- 412. ** K. Abe *et al.* [BELLE Collaboration], "Improved measurement of mixing-induced CP violation in the neutral *B* meson system," Phys. Rev. D **66**, 071102(R) (2002).
- 413. T. Affolder et al. [CDF Collaboration], "Diffractive dijet production at $\sqrt{s} = 630$ -GeV and 1800-GeV at the Fermilab Tevatron," Phys. Rev. Lett. 88, 151802 (2002).
- 414. D. Acosta *et al.* [CDF Collaboration], "Search for new physics in photon lepton events in proton antiproton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **66**, 012004 (2002)
- 415. D. Acosta *et al.* [CDF Collaboration], "Search for the decay $B_s^0 \to \mu^+\mu^-\phi$ in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **65**, 111101 (2002).
- 416. D. Acosta *et al.* [CDF Collaboration], 'Search for single-top-quark production in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **65**, 091102 (2002)

- 417. D. Acosta *et al.* [CDF Collaboration], 'Comparison of the isolated direct photon cross sections in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV and $\sqrt{s} = 0.63$ -TeV," Phys. Rev. D **65**, 112003 (2002)
- 418. ** K. Abe *et al.* [Belle Collaboration], "Observation of mixing-induced CP violation in the neutral B meson system," Phys. Rev. D **66**, 032007 (2002)
- 419. D. Acosta *et al.* [CDF Collaboration], "Search for new physics in photon lepton events in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **89**, 041802 (2002)
- 420. ** K. Abe *et al.* [BELLE Collaboration], "Measurement of CP-violating asymmetries in $B0 \to \pi^+ \pi^-$ decays," Phys. Rev. Lett. **89**, 071801 (2002).
- 421. D. Acosta *et al.* [CDF Collaboration], "Upsilon production and polarization in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **88**, 161802 (2002).
- 422. D. Acosta *et al.* [CDF Collaboration], "Soft and hard interactions in $p \bar{p}$ collisions at \sqrt{s} = 1800-GeV and 630-GeV," Phys. Rev. D **65**, 072005 (2002).
- 423. ** K. Abe *et al.* [BELLE Collaboration], "Production of prompt charmonia in e^+ e^- annihilation at $\sqrt{s} = 10.6$ -GeV," Phys. Rev. Lett. **88**, 052001 (2002).
- 424. D. Acosta *et al.* [CDF Collaboration], "Study of the heavy flavor content of jets produced in association with W bosons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **65**, 052007 (2002).
- 425. T. Affolder *et al.* [CDF Collaboration], 'Measurement of the strong coupling constant from inclusive jet production at the Tevatron \bar{p} p collider," Phys. Rev. Lett. **88**, 042001 (2002).
- 426. T. Affolder *et al.* [CDF Collaboration], "A Study of $B^0 \to J/\psi K^{(*)0}\pi^+\pi^-$ decays with the Collider Detector at Fermilab," Phys. Rev. Lett. **88**, 071801 (2002).
- 427. ** W. Adam *et al.* [RD42 Collaboration], "Diamond Pixel Detectors," Nucl. Instrum. Meth. A **465**, 88 (2000).
- 428. T. Affolder *et al.* [CDF Collaboration], "Search for new heavy particles in the WZ^0 final state in $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV," Phys. Rev. Lett. **88**, 071806 (2002).
- 429. T. Affolder *et al.* [CDF Collaboration], "Observation of diffractive J/ψ production at the Fermilab Tevatron," Phys. Rev. Lett. **87**, 241802 (2001).
- 430. T. Affolder *et al.* [CDF Collaboration], "Double diffraction dissociation at the Fermilab Tevatron collider," Phys. Rev. Lett. **87**, 141802 (2001).
- 431. T. Affolder *et al.* [CDF Collaboration], "Charged particle multiplicity in jets in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **87**, 211804 (2001).

- 432. T. Affolder *et al.* [CDF Collaboration], "Search for quark lepton compositeness and a heavy W' boson using the $e\nu$ channel in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. 87, 231803 (2001).
- 433. T. Affolder *et al.* [CDF Collaboration], "Searches for new physics in events with a photon and *b*-quark jet at CDF," Phys. Rev. D **65**, 052006 (2002).
- 434. T. Affolder *et al.* [CDF Collaboration], "Search for gluinos and squarks using like-sign dileptons in $p \bar{p}$ collisions $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **87**, 251803 (2001).
- 435. T. Affolder *et al.* [CDF Collaboration], "Measurement of $d\sigma/dM$ forward-backward charge asymmetry for high mass Drell-Yan e^+ e^- pairs from p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **87**, 131802 (2001).
- 436. T. Affolder *et al.* [CDF Collaboration], "Cross section and heavy quark composition of $\gamma + \mu$ events produced in $p \bar{p}$ collisions," Phys. Rev. D **65**, 012003 (2002).
- 437. T. Affolder *et al.* [CDF Collaboration], "Search for gluinos and scalar quarks in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV using the missing energy plus multijets signature," Phys. Rev. Lett. **88**, 041801 (2002).
- 438. T. Affolder *et al.* [CDF Collaboration], "Production of χ_{c1} and χ_{c2} in $p \bar{p}$ collisions at \sqrt{s} = 1.8-TeV," Phys. Rev. Lett. **86**, 3963 (2001).
- 439. T. Affolder *et al.* [CDF Collaboration], "Search for narrow diphoton resonances and for $\gamma \gamma + W/Z$ signatures in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **64**, 092002 (2001).
- 440. T. Affolder *et al.* [CDF Collaboration], "Measurement of the inclusive jet cross section in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **64**, 032001 (2001). [Erratum-ibid. D **65**, 039903 (2001)]
- 441. T. Affolder *et al.* [CDF Collaboration], "Measurement of the $t\bar{t}$ production cross section in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **64**, 032002 (2001).
- 442. T. Affolder *et al.* [CDF Collaboration], "First measurement of the ratio $\mathcal{B}(t \to Wb)/\mathcal{B}(t \to Wq)$ and associated limit on the CKM element |V(tb)|," Phys. Rev. Lett. **86**, 3233 (2001).
- 443. T. Affolder *et al.* [CDF Collaboration], "Measurement of the two-jet differential cross section in proton antiproton collisions at $\sqrt{s} = 1800$ -GeV," Phys. Rev. D **64**, 012001 (2001). [Erratum-ibid. D **65**, 039902 (2001)]
- 444. T. Affolder *et al.* [CDF Collaboration], "Measurement of the top quark p_T distribution," Phys. Rev. Lett. **87**, 102001 (2001).
- 445. T. Affolder *et al.* [CDF Collaboration], "Search for the supersymmetric partner of the top quark in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **63**, 091101 (2001).
- 446. T. Affolder *et al.* [CDF Collaboration], "Search for neutral supersymmetric Higgs bosons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **86**, 4472 (2001).

- 447. T. Affolder *et al.* [CDF Collaboration], "Test of enhanced leading order QCD in W boson plus jets events from 1.8-TeV \bar{p} p collisions," Phys. Rev. D **63**, 072003 (2001).
- 448. W. Adam *et al.* [RD42 Collaboration], "Performance of irradiated CVD diamond microstrip sensors," Nucl. Instrum. Meth. A **476**, 706 (2002).
- 449. W. Adam *et al.* [RD42 Collaboration], "Pulse height distribution and radiation tolerance of CVD Diamond detectors," Nucl. Instrum. Meth. A **447**, 244 (2000).
- 450. F. Hartjes *et al.* [RD42 Collaboration], "Parameterisation Of radiation effects on CVD Diamond for proton irradiation," Nucl. Phys. Proc. Suppl. **78**, 675 (1999).
- 451. R. Wedenig *et al.* [RD42 Collaboration], "CVD Diamond pixel detectors for LHC experiments," Nucl. Phys. Proc. Suppl. **78**, 497 (1999).
- 452. W. Adam *et al.* [RD42 Collaboration], "Recent results with CVD Diamond trackers," Nucl. Phys. Proc. Suppl. **78**, 329 (1999).
- 453. J. Armitage *et al.* [ATLAS-Canada Collaboration], "Construction and initial beam tests of the ATLAS Tungsten forward calorimeter," Nucl. Phys. Proc. Suppl. **78**, 171 (1999).
- 454. T. Affolder *et al.* [CDF Collaboration], "Measurement of the W boson mass with the Collider Detector at Fermilab," Phys. Rev. D **64**, 052001 (2001).
- 455. T. Affolder *et al.* [CDF Collaboration], "Measurement of the decay amplitudes of $B^0 \to J/\psi K^{*0}$ and $B_s^0 \to J/\psi \phi$ decays," Phys. Rev. Lett. **85**, 4668 (2000).
- 456. T. Affolder *et al.* [CDF Collaboration], "Measurement of the top quark mass with the Collider Detector at Fermilab," Phys. Rev. D **63**, 032003 (2001).
- 457. T. Affolder *et al.* [CDF Collaboration], "Measurement of $d\sigma/dy$ for high mass Drell-Yan e^+e^- pairs from p \bar{p} collisions at $\sqrt{s}=1.8$ -TeV," Phys. Rev. D **63**, 011101 (2001).
- 458. T. Affolder *et al.* [CDF Collaboration], "Dijet production by double pomeron exchange at the Fermilab Tevatron," Phys. Rev. Lett. **85**, 4215 (2000).
- 459. ** T. Affolder *et al.* [CDF Collaboration], "Measurement of J/ψ and $\psi(2S)$ polarization in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **85**, 2886 (2000).
- 460. T. Affolder *et al.* [CDF Collaboration], "Search For color singlet technicolor particles in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 1110 (2000).
- 461. T. Affolder et al. [CDF Collaboration], "Direct measurement of the W boson width in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **85**, 3347 (2000).
- 462. W. Adam *et al.* [RD42 Collaboration], "Micro-strip sensors based on CVD diamond," Nucl. Instrum. Meth. A **453**, 141 (2000).
- 463. T. Affolder *et al.* [CDF Collaboration], "Search for second and third generation leptoquarks including production via technicolor interactions in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **85**, 2056 (2000).

- 464. T. Affolder *et al.* [CDF Collaboration], "Limits on gravitino production and new processes with large missing transverse energy in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **85**, 1378 (2000).
- 465. T. Affolder *et al.* [CDF Collaboration], "Diffractive dijets with a leading antiproton in \bar{p} p collisions at $\sqrt{s} = 1800$ -GeV," Phys. Rev. Lett. **84**, 5043 (2000).
- 466. T. Affolder *et al.* [CDF Collaboration], "Search for new particles decaying to $t\bar{t}$ in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **85**, 2062 (2000).
- 467. M. Friedl *et al.* [RD42 Collaboration], "CVD Diamond detectors for ionizing radiation," Nucl. Instrum. Meth. A **435**, 194 (1999).
- 468. ** W. Adam *et al.* [RD42 Collaboration], "The first bump bonded pixel detectors on CVD Diamond," Nucl. Instrum. Meth. A **436**, 326 (1999).
- 469. ** T. Affolder *et al.* [CDF Collaboration], "A measurement of the differential dijet mass cross section in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **61**, 091101 (2000).
- 470. T. Affolder *et al.* [CDF Collaboration], "Search for scalar top quark production in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 5273 (2000).
- 471. T. Affolder *et al.* [CDF Collaboration], "Observation of orbitally excited *B* mesons in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **64**, 072002 (2001).
- 472. T. Affolder *et al.* [CDF Collaboration], "Search for the charged Higgs boson in the decays of top quark pairs in the $e\tau$ and $\mu\tau$ channels at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **62**, 012004 (2000).
- 473. W. Adam *et al.* [RD42 Collaboration], "Review of the development of Diamond radiation sensors," Nucl. Instrum. Meth. A **434**, 131 (1999).
- 474. T. Affolder *et al.* [CDF Collaboration], "Search for scalar top and scalar bottom quarks in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 5704 (2000).
- 475. T. Affolder *et al.* [CDF Collaboration], "Production of Upsilon(1S) mesons from χ_b decays in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 2094 (2000).
- 476. ** F. Abe *et al.* [CDF Collaboration], "Measurement of *b* quark fragmentation fractions in the production of strange and light *B* mesons in $p \bar{p}$ collisions at \sqrt{s} =1.8-TeV," Phys. Rev. D **60**, 092005 (1999).
- 477. F. Abe et al. [CDF Collaboration], "Search for a W' boson via the decay mode $W' \to \mu\nu_{\mu}$ in 1.8-TeV $p \bar{p}$ collisions," Phys. Rev. Lett. 84, 5716 (2000).
- 478. T. Affolder *et al.* [CDF Collaboration], "Measurement of the helicity of W bosons in top quark decays," Phys. Rev. Lett. **84**, 216 (2000).
- 479. S. Schnetzer *et al.*, "Tracking with CVD Diamond radiation sensors at high luminosity colliders," IEEE Trans. Nucl. Sci. **46**, 193 (1999).

- 480. T. Affolder *et al.* [CDF Collaboration], "Search for a fourth-generation quark more massive than the Z^0 boson in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 835 (2000).
- 481. T. Affolder *et al.* [CDF Collaboration], "Measurement of *b* quark fragmentation fractions in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **84**, 1663 (2000).
- 482. T. Affolder *et al.* [CDF Collaboration], "A measurement of $\sin(2\beta)$ from $B \to J/\psi K_S^0$ with the CDF detector," Phys. Rev. D **61**, 072005 (2000).
- 483. T. Affolder *et al.* [CDF Collaboration], "The transverse momentum and total cross section of e^+ e^- pairs in the Z boson region from p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. 84, 845 (2000).
- 484. T. Affolder *et al.* [CDF Collaboration], "Observation of diffractive beauty production at the Fermilab Tevatron," Phys. Rev. Lett. **84**, 232 (2000).
- 485. T. Affolder *et al.* [CDF Collaboration], "Measurement of the $B^0\bar{B}^0$ oscillation frequency using l^-D^{*+} pairs and lepton flavor tags," Phys. Rev. D **60**, 112004 (1999).
- 486. T. Affolder *et al.* [CDF Collaboration], "Search for the flavor-changing neutral current decays $B^+ \to \mu^+ \mu^- K^+$ and $B^0 \to \mu^+ \mu^- K^{*0}$," Phys. Rev. Lett. **83**, 3378 (1999).
- 487. C. Bauer *et al.*, "Recent results from the RD42 Diamond detector collaboration," Nucl. Instrum. Meth. A **383**, 64 (1996).
- 488. F. Abe *et al.* [The CDF Collaboration], "Measurement of the $B^0\bar{B}^0$ oscillation frequency using dimuon data in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **60**, 051101 (1999).
- 489. F. Abe *et al.* [CDF Collaboration], "Measurement of $B^0\bar{B}^0$ flavor oscillations using jet-charge and lepton flavor tagging in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **60**, 072003 (1999).
- 490. F. Abe *et al.* [CDF Collaboration], "A search for $B_s^0 \bar{B}_s^0$ oscillations using the semileptonic decay $B_s^0 \to \phi l^+ X \nu$," Phys. Rev. Lett. **82**, 3576 (1999).
- 491. C. Bauer *et al.* [RD42 Collaboration], "Radiation hardness studies of CVD Diamond detectors," Nucl. Instrum. Meth. A **367**, 207 (1995).
- 492. C. Bauer *et al.* [RD42 Collaboration], "Recent results from Diamond microstrip detectors," Nucl. Instrum. Meth. A **367**, 202 (1995).
- 493. F. Abe *et al.* [CDF Collaboration], "Measurement of $b\bar{b}$ rapidity correlations in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **61**, 032001 (2000).
- 494. F. Abe *et al.* [CDF Collaboration], "Search for R-parity violating supersymmetry using like-sign dielectrons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **83**, 2133 (1999).
- 495. F. Abe *et al.* [CDF Collaboration], "Events with a rapidity gap between jets in \bar{p} p collisions at $\sqrt{s} = 630$ -GeV," Phys. Rev. Lett. **81**, 5278 (1998).

- 496. F. Abe *et al.* [CDF Collaboration], "Search for third-generation leptoquarks from technicolor models in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **82**, 3206 (1999).
- 497. F. Abe *et al.* [CDF Collaboration], "Kinematics of $t\bar{t}$ events at CDF," Phys. Rev. D **59**, 092001 (1999).
- 498. F. Abe *et al.* [CDF Collaboration], "Search for a technicolor Ω_T particle in events with a photon and a *b* quark jet at CDF," Phys. Rev. Lett. **83**, 3124 (1999).
- 499. F. Abe *et al.* [CDF Collaboration], "Measurement of the top quark mass with the Collider Detector at Fermilab," Phys. Rev. Lett. **82**, 271 (1999). [Erratum-ibid. **82**, 2808 (1999)]
- 500. P. Weilhammer *et al.* [RD42 Collaboration], "Recent results on CVD diamond radiation sensors," Nucl. Instrum. Meth. A **409**, 264 (1998).
- 501. F. Abe *et al.* [CDF Collaboration], "Measurement of Z^0 and Drell-Yan production cross section using dimuons in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **59**, 052002 (1999).
- 502. F. Abe *et al.* [CDF Collaboration], "Search for new particles decaying to $b\bar{b}$ in $p\ \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **82**, 2038 (1999).
- 503. F. Abe *et al.* [CDF Collaboration], "Search for the decays $B^0, B_s^0 \to e^{+/-} \mu^{-/+}$ and Pati-Salam leptoquarks," Phys. Rev. Lett. **81**, 5742 (1998).
- 504. F. Abe *et al.* [CDF Collaboration], "Measurement of the lepton charge asymmetry in W boson decays produced in $p \bar{p}$ collisions," Phys. Rev. Lett. **81**, 5754 (1998).
- 505. F. Abe *et al.* [CDF Collaboration], "Search for Higgs bosons produced in association with a vector boson in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **81**, 5748 (1998).
- 506. M. Zoeller *et al.* [RD42 Collaboration], "Performance of CVD Diamond microstrip detectors under particle irradiation," IEEE Trans. Nucl. Sci. **44**, 815 (1997).
- 507. F. Abe *et al.* [CDF Collaboration], "Measurement of the B_s^0 meson lifetime using semileptonic decays," Phys. Rev. D **59**, 032004 (1999).
- 508. F. Abe *et al.* [CDF Collaboration], "Search for second generation leptoquarks in the dimuon plus dijet channel of $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **81**, 4806 (1998).
- 509. F. Abe *et al.* [CDF Collaboration], "Searches for new physics in diphoton events in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **59**, 092002 (1999).
- 510. F. Abe *et al.* [CDF Collaboration], "Measurement of the $B^0\bar{B}^0$ flavor oscillation frequency and study of same side flavor tagging of B mesons in p \bar{p} collisions," Phys. Rev. D **59**, 032001 (1999).
- 511. F. Abe *et al.* [CDF Collaboration], "Measurement of the CP-violation parameter $\sin(2\beta)$ in $B^0, \bar{B}^0 \to J/\psi K_S^0$ decays," Phys. Rev. Lett. **81**, 5513 (1998).

- 512. F. Abe *et al.* [CDF Collaboration], "Improved measurement of the B^- and \bar{B}^0 meson lifetimes using semileptonic decays," Phys. Rev. D **58**, 092002 (1998).
- 513. D. Meier *et al.* [RD42 Collaboration], "Proton irradiation of CVD diamond detectors for high-luminosity experiments at the LHC," Nucl. Instrum. Meth. A **426**, 173 (1999).
- 514. F. Abe *et al.* [CDF Collaboration], "Observation of the B_c^+ meson in $p \bar{p}$ collisions at \sqrt{s} = 1.8-TeV," Phys. Rev. Lett. **81**, 2432 (1998).
- 515. F. Abe *et al.* [The CDF Collaboration], "Search for long-lived parents of Z^0 bosons in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **58**, 051102 (1998).
- 516. F. Abe *et al.* [CDF Collaborations], "Measurement of the $\sigma(W + \ge 1 \text{ jet})/\sigma(W)$ cross section ratio from \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **81**, 1367 (1998).
- 517. F. Abe *et al.* [CDF Collaboration], "Observation of B_c^+ mesons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **58**, 112004 (1998).
- 518. F. Abe et al. [CDF Collaboration], "Search for the rare decay $W^{\pm} \to D_s^{\pm} \gamma$ in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D 58, 091101 (1998).
- 519. F. Abe *et al.* [CDF Collaboration], "Search For the rare Decay $W^{\pm} \to \pi^{\pm} + \gamma$ in proton antiproton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **58**, 031101 (1998).
- 520. ** F. Abe *et al.* [CDF Collaboration], "Observation of $B^+ \to \psi(2S)K^+$ and $B^0 \to \psi(2S)K^{*0}(892)$ decays and measurements of B meson branching fractions into J/ψ and $\psi(2S)$ final states," Phys. Rev. D **58**, 072001 (1998).
- 521. F. Abe *et al.* [CDF collaboration], "Search for chargino neutralino associated production at the Fermilab Tevatron Collider," Phys. Rev. Lett. **80**, 5275 (1998).
- 522. F. Abe *et al.* [CDF Collaboration], "Searches for new physics in diphoton events in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **81**, 1791 (1998).
- 523. F. Abe *et al.* [CDF Collaboration], "Jet pseudorapidity distribution in direct photon events in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **57**, 1359 (1998).
- 524. F. Abe *et al.* [CDF Collaboration], "Search for the decays $B^0 \to \mu^+ \mu^-$ and $B_s^0 \to \mu^+ \mu^-$ in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **57**, 3811 (1998).
- 525. F. Abe *et al.* [CDF Collaboration], "Observation of hadronic W decays in $t\bar{t}$ events with the Collider Detector at Fermilab," Phys. Rev. Lett. **80**, 5720 (1998).
- 526. W. Adam *et al.* [RD42 Collaboration], "Status of diamond particle detectors," Nucl. Instrum. Meth. A **418**, 197 (1998).
- 527. F. Abe *et al.* [CDF Collaboration], "Measurement of B hadron lifetimes using J/ψ final states at CDF," Phys. Rev. D **57**, 5382 (1998).

- 528. F. Abe *et al.* [CDF Collaboration], "Measurement of the top quark mass and $t\bar{t}$ production cross section from dilepton events at the Collider Detector at Fermilab," Phys. Rev. Lett. **80**, 2779 (1998).
- 529. F. Abe *et al.* [CDF Collaboration], "Measurement of the top quark mass," Phys. Rev. Lett. **80**, 2767 (1998).
- 530. F. Abe *et al.* [CDF Collaboration], "Measurement of the $t\bar{t}$ production cross section in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **80**, 2773 (1998).
- 531. F. Abe *et al.* [CDF Collaboration], "Measurement of the $B^0\bar{B}^0$ oscillation frequency using πB meson charge-flavor correlations in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **80**, 2057 (1998).
- 532. F. Abe *et al.* [CDF Collaboration], "Search for flavor-changing neutral current decays of the top quark in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **80**, 2525 (1998).
- 533. F. Abe *et al.* [CDF Collaboration], "Properties of jets in W boson events from 1.8-TeV \bar{p} p collisions," Phys. Rev. Lett. **79**, 4760 (1997).
- 534. F. Abe *et al.* [CDF Collaboration], "Measurement of the differential cross section for events with large total transverse energy in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **80**, 3461 (1998).
- 535. F. Abe *et al.* [CDF Collaboration], "Dijet production by color-singlet exchange at the Fermilab Tevatron," Phys. Rev. Lett. **80**, 1156 (1998).
- 536. F. Abe *et al.* [CDF Collaboration], "Search for first generation leptoquark pair production in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 4327 (1997).
- 537. F. Abe *et al.* [CDF Collaboration], "Search for new particles decaying into $b\bar{b}$ and produced in association with W bosons decaying into $e \nu$ or $\mu \nu$ at the Tevatron," Phys. Rev. Lett. **79**, 3819 (1997).
- 538. F. Abe *et al.* [CDF Collaboration], "Limits on quark-lepton compositeness scales from dileptons produced in 1.8-TeV $p \bar{p}$ collisions," Phys. Rev. Lett. **79**, 2198 (1997).
- 539. F. Abe *et al.* [CDF Collaboration], "Search for new gauge bosons decaying into dileptons in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 2192 (1997).
- 540. D. Husson *et al.*, "Neutron irradiation of CVD diamond samples for tracking detectors," Nucl. Instrum. Meth. A **388**, 421 (1997).
- 541. F. Abe *et al.* [CDF Collaboration], "Properties of photon plus two-jet events in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **57**, 67 (1998).
- 542. F. Abe *et al.* [CDF Collaboration], "Properties of six-jet events with large six-jet mass at the Fermilab proton anti-proton collider," Phys. Rev. D **56**, 2532 (1997).

- 543. F. Abe *et al.* [CDF Collaboration], "Double parton scattering in \bar{p} p collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **56**, 3811 (1997).
- 544. F. Abe et al. [CDF Collaboration], "The $\mu\tau$ and $e\tau$ decays of top quark pairs produced in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 3585 (1997).
- 545. F. Abe *et al.* [CDF Collaboration], "Measurement of double parton scattering in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 584 (1997).
- 546. F. Abe *et al.* [CDF Collaboration], "Search for charged Higgs decays of the top quark using hadronic decays of the tau lepton," Phys. Rev. Lett. **79**, 357 (1997).
- 547. F. Abe *et al.* [CDF Collaboration], "First observation of the all hadronic decay of $t\bar{t}$ pairs," Phys. Rev. Lett. **79**, 1992 (1997).
- 548. F. Abe *et al.* [CDF Collaboration], "Measurement of diffractive dijet production at the Tevatron," Phys. Rev. Lett. **79**, 2636 (1997).
- 549. D. Husson *et al.* [RD42 Collaboration], "Neutron irradiation of CVD Diamond samples for tracking detectors," Nucl. Instrum. Meth. A **388**, 421 (1997).
- 550. F. Abe *et al.* [CDF Collaboration], "Production of J/ψ mesons from χ_c meson decays in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 578 (1997).
- 551. F. Abe *et al.* [CDF Collaboration], "Search for gluinos and squarks at the Fermilab Tevatron collider," Phys. Rev. D **56**, 1357 (1997).
- 552. F. Abe *et al.* [CDF Collaboration], " J/ψ and $\psi(2S)$ production in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **79**, 572 (1997).
- 553. F. Abe *et al.* [CDF Collaboration], "Observation of diffractive W boson production at the Tevatron," Phys. Rev. Lett. **78**, 2698 (1997).
- 554. F. Abe *et al.* [CDF Collaboration], "Search for new particles decaying to dijets at CDF," Phys. Rev. D **55**, 5263 (1997).
- 555. ** P. Abreu *et al.* [DELPHI Collaboration], "A precise measurement of the B⁰ meson lifetime using a new technique," Z. Phys. C **74**, 19 (1997) [Erratum-ibid. C **75**, 579 (1997)].
- 556. W. Adam *et al.* [DELPHI Collaboration], "Search for pair production of heavy objects in 4-jet events at $\sqrt{s} = 130$ -GeV to 136-GeV," Z. Phys. C **73**, 1 (1996).
- 557. P. Abreu *et al.* [DELPHI Collaboration], "An upper limit for $\mathcal{B}(Z^0 \to ggg)$ from symmetric 3-jet Z^0 hadronic decays," Phys. Lett. B **389**, 405 (1996).
- 558. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of event shape and inclusive distributions at $\sqrt{s} = 130$ -GeV and 136-GeV," Z. Phys. C **73**, 229 (1997).

- 559. P. Abreu *et al.* [DELPHI Collaboration], "Search for lepton flavour number violating Z^0 decays," Z. Phys. C **73**, 243 (1997).
- 560. P. Abreu *et al.* [DELPHI Collaboration], "Tuning and test of fragmentation models based on identified particles and precision event shape data," Z. Phys. C **73**, 11 (1996).
- 561. P. Abreu *et al.* [DELPHI Collaboration], "Search for neutralinos, scalar leptons and scalar quarks in e^+ e^- interactions at $\sqrt{s} = 130$ -GeV and 136-GeV," Phys. Lett. B **387**, 651 (1996).
- 562. W. Adam *et al.* [DELPHI Collaboration], "Study of rare *b* decays with the DELPHI detector at LEP," Z. Phys. C **72**, 207 (1996).
- 563. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of inclusive $K^{*0}(892)$, ϕ and $K_2^{*0}(1430)$ production in hadronic Z decays," Z. Phys. C **73**, 61 (1996).
- 564. P. Abreu *et al.* [DELPHI Collaboration], "Search for high mass $\gamma\gamma$ resonances in $e^+e^- \to l^+l^-\gamma\gamma$, $\nu\bar{\nu}\gamma\gamma$ and $q\bar{q}\gamma\gamma$ at LEP I," Z. Phys. C **72**, 179 (1996).
- 565. P. Abreu *et al.* [DELPHI Collaboration], "Search for the lightest chargino at $\sqrt{s} = 130$ -GeV and 136-GeV in DELPHI," Phys. Lett. B **382**, 323 (1996).
- 566. P. Abreu *et al.* [DELPHI Collaboration], "Study of radiative leptonic events with hard photons and search for excited charged leptons at $\sqrt{s} = 130$ -GeV to 136-GeV," Phys. Lett. B **380**, 480 (1996).
- 567. W. Adam *et al.* [DELPHI Collaboration], "Search for anomalous production of single photons at $\sqrt{s} = 130$ -GeV and 136-GeV," Phys. Lett. B **380**, 471 (1996).
- 568. P. Abreu *et al.* [DELPHI Collaboration], "Kaon interference in the hadronic decays of the Z^0 ," Phys. Lett. B **379**, 330 (1996).
- 569. P. Abreu *et al.* [DELPHI Collaboration], "First study of the interference between initial and final state radiation at the Z resonance," Z. Phys. C **72**, 31 (1996).
- 570. P. Abreu *et al.* [DELPHI Collaboration], "Mean lifetime of the B_s^0 meson," Z. Phys. C **71**, 11 (1996).
- 571. P. Abreu *et al.* [DELPHI Collaboration], "First measurement of $f'_2(1525)$ production in Z^0 hadronic decays," Phys. Lett. B **379**, 309 (1996).
- 572. P. Abreu *et al.* [DELPHI Collaboration], "Determination of the average lifetime of b baryons," Z. Phys. C **71**, 199 (1996).
- 573. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the partial decay width $R(b) = \Gamma(b\bar{b}) / \Gamma(had)$ of the Z with the DELPHI detector at LEP," Z. Phys. C **70**, 531 (1996).
- 574. P. Abreu *et al.* [DELPHI Collaboration], "Search for exclusive decays of the Λ_b baryon and measurement of its mass," Phys. Lett. B **374**, 351 (1996).

- 575. P. Abreu *et al.* [DELPHI Collaboration], "Determination of |V(cb)| from the semileptonic decay $B^0 \to D^{*-}$ lepton neutrino," Z. Phys. C **71**, 539 (1996).
- 576. W. Adam *et al.* [DELPHI Collaboration], "Production of Σ_0 and Ω^- in Z decays," Z. Phys. C **70**, 371 (1996).
- 577. P. Abreu *et al.* [DELPHI Collaboration], "Updated precision measurement of the average lifetime of *B* hadrons," Phys. Lett. B **377**, 195 (1996).
- 578. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the B^0 oscillation frequency using kaons, leptons and jet charge," Z. Phys. C **72**, 17 (1996).
- 579. P. Abreu *et al.* [DELPHI Collaboration], "Search for new phenomena using single photon events in the DELPHI detector at LEP," Z. Phys. C **74**, 577 (1997).
- 580. P. Abreu *et al.* [DELPHI Collaboration], "Charged particle multiplicity in e^+ e^- interactions at $\sqrt{s} = 130$ -GeV," Phys. Lett. B **372**, 172 (1996).
- 581. ** P. Abreu *et al.* [DELPHI Collaboration], "Performance of the DELPHI detector," Nucl. Instrum. Meth. A **378**, 57 (1996).
- 582. P. Abreu *et al.* [DELPHI Collaboration], "Energy dependence of the differences between the quark and gluon jet fragmentation," Z. Phys. C **70**, 179 (1996).
- 583. ** P. Abreu *et al.* [DELPHI Collaboration], "A Precise measurement of the tau lepton lifetime," Phys. Lett. B **365**, 448 (1996).
- 584. P. Abreu *et al.* [DELPHI Collaboration], "Search for promptly produced heavy quarkonium states in hadronic Z decays," Z. Phys. C **69**, 575 (1996).
- 585. W. Adam et al. [DELPHI Collaboration], "Measurement of inclusive π^0 production in hadronic Z^0 decays," Z. Phys. C **69**, 561 (1996).
- 586. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of $\Delta^{++}(1232)$ production in hadronic Z decays," Phys. Lett. B **361**, 207 (1995).
- 587. P. Abreu *et al.* [DELPHI Collaboration], "Upper limits on the branching ratios $\tau \to \mu \gamma$ and $\tau \to e \gamma$," Phys. Lett. B **359**, 411 (1995).
- 588. ** P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the tau leptonic branching fractions," Phys. Lett. B **357**, 715 (1995).
- 589. P. Abreu *et al.* [DELPHI Collaboration], "Study of prompt photon production in hadronic Z^0 decays," Z. Phys. C **69**, 1 (1995).
- 590. P. Abreu *et al.* [DELPHI Collaboration], "Search for exclusive charmless B meson decays with the DELPHI detector at LEP," Phys. Lett. B **357**, 255 (1995).
- 591. ** V. Chabaud *et al.* [DELPHI Collaboration], "The DELPHI silicon strip microvertex detector with double sided readout," Nucl. Instrum. Meth. A **368**, 314 (1996).

- 592. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the photon structure function F2(gamma) at an average q^2 of 12-GeV²/ c^4 ," Z. Phys. C **69**, 223 (1996).
- 593. P. Abreu *et al.* [DELPHI Collaboration], "Observation of short range three particle correlations in e^+ e^- annihilations at LEP energies," Phys. Lett. B **355**, 415 (1995).
- 594. P. Abreu *et al.* [DELPHI Collaboration], "Lifetime and production rate of beauty baryons from Z decays," Z. Phys. C **68**, 375 (1995).
- 595. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of B^+ and B^0 lifetimes using $\bar{D}l^+$ events," Z. Phys. C **68**, 13 (1995).
- 596. W. Adam *et al.* [DELPHI Collaboration], "Lifetimes of charged and neutral *B* hadrons using event topology," Z. Phys. C **68**, 363 (1995).
- 597. P. Abreu *et al.* [DELPHI Collaboration], " B^* production in Z decays," Z. Phys. C **68**, 353 (1995).
- 598. P. Abreu *et al.* [DELPHI Collaboration], "Strange baryon production in Z hadronic decays," Z. Phys. C **67**, 543 (1995).
- 599. P. Abreu et al. [DELPHI Collaboration], "Inclusive measurements of the K^{\pm} and p / \bar{p} production in hadronic Z^0 decays," Nucl. Phys. B **444**, 3 (1995).
- 600. P. Abreu et al. [DELPHI Collaboration], "Measurements of the tau polarization in \mathbb{Z}^0 decays," Z. Phys. C **67**, 183 (1995).
- 601. P. Abreu *et al.* [DELPHI Collaboration], "Production of strange B baryons decaying into Ξ^{\pm} lepton pairs at LEP," Z. Phys. C **68**, 541 (1995).
- 602. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of $\Gamma(b\bar{b})/\Gamma(\text{hadrons})$ using impact parameter measurements and lepton identification," Z. Phys. C **66**, 323 (1995).
- 603. F. Abe *et al.* [CDF Collaboration], "Measurement of the *B* meson and *b* quark cross-sections at $\sqrt{s} = 1.8$ -TeV using the exclusive decay $B^0 \to J/\psi K^{*0}(892)$," Phys. Rev. D **50**, 4252 (1994).
- 604. P. Abreu *et al.* [DELPHI Collaboration], "Production of charged particles, K_S^0 , K^{\pm} , p and Λ in $Z \to b\bar{b}$ events and in the decay of b hadrons," Phys. Lett. B **347**, 447 (1995).
- 605. P. Abreu *et al.* [DELPHI Collaboration], "First measurement of the strange quark asymmetry at the Z^0 peak," Z. Phys. C **67**, 1 (1995).
- 606. P. Abreu *et al.* [DELPHI Collaboration], "Search for heavy neutral Higgs bosons in two doublet models," Z. Phys. C **67**, 69 (1995).
- 607. P. Abreu *et al.* [DELPHI Collaboration], "Observation of orbitally excited *B* mesons," Phys. Lett. B **345**, 598 (1995).

- 608. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the forward backward asymmetry of charm and bottom quarks at the Z pole using $D^{*\pm}$ mesons," Z. Phys. C **66**, 341 (1995).
- 609. P. Abreu *et al.* [DELPHI Collaboration], "First evidence of hard scattering processes in single tagged gamma gamma collisions," Phys. Lett. B **342**, 402 (1995).
- 610. P. Abreu et al. [DELPHI Collaboration], "Measurement of the forward backward asymmetry of $e^+ e^- \to Z \to b\bar{b}$ using prompt leptons and a lifetime tag," Z. Phys. C 65, 569 (1995).
- 611. ** F. Borchelt *et al.*, "First measurements with a diamond microstrip detector," Nucl. Instrum. Meth. A **354**, 318 (1995).
- 612. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of time dependent $B^0\bar{B}^0$ mixing," Phys. Lett. B **338**, 409 (1994).
- 613. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the $\Gamma(b\bar{b})/\Gamma(\text{hadron})$ branching ratio of the Z by double hemisphere tagging," Z. Phys. C **65**, 555 (1995).
- 614. P. Abreu *et al.* [DELPHI Collaboration], Z. Phys. C **65**, 587 (1995).
- 615. P. Abreu *et al.* [DELPHI Collaboration], " J/ψ production in the hadronic decays of the Z," Phys. Lett. B **341**, 109 (1994).
- 616. P. Abreu *et al.* [DELPHI Collaboration], "A Study of radiative muon pair events at Z^0 energies and limits on an additional Z' gauge boson," Z. Phys. C **65**, 603 (1995).
- 617. P. Abreu *et al.* [DELPHI Collaboration], "Charged kaon production in tau decays at LEP," Phys. Lett. B **334**, 435 (1994).
- 618. P. Abreu *et al.* [DELPHI Collaboration], "Search for pair produced heavy scalars in \mathbb{Z}^0 decays," Z. Phys. C **64**, 183 (1994).
- 619. F. Abe *et al.* [CDF Collaboration], "Search for radiative decays of neutralinos in proton anti-proton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **75**, 613 (1995).
- 620. F. Abe *et al.* [CDF Collaboration], "Observation of rapidity gaps in \bar{p} p collisions at 1.8-TeV," Phys. Rev. Lett. **74**, 855 (1995).
- 621. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the $B\bar{B}$ mixing parameter in DELPHI," Phys. Lett. B **332**, 488 (1994).
- 622. P. Abreu *et al.* [DELPHI Collaboration], "Search for the standard model Higgs boson in Z^0 decays," Nucl. Phys. B **421**, 3 (1994).
- 623. F. Abe *et al.* [CDF Collaboration], "Evidence for color coherence in $p \bar{p}$ collisions at \sqrt{s} = 1.8-TeV," Phys. Rev. D **50**, 5562 (1994).

- 624. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the $e^+e^- \to \gamma\gamma(\gamma)$ cross-section at LEP energies," Phys. Lett. B **327**, 386 (1994).
- 625. P. Abreu *et al.* [DELPHI Collaboration], "Improved measurements of cross-sections and asymmetries at the Z^0 resonance," Nucl. Phys. B **418**, 403 (1994).
- 626. ** P. Abreu *et al.* [DELPHI Collaboration], "A Precision measurement of the average lifetime of *B* hadrons," Z. Phys. C **63**, 3 (1994).
- 627. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the B_s^0 meson mass," Phys. Lett. B **324**, 500 (1994).
- 628. P. Abreu *et al.* [DELPHI Collaboration], "Measurements of the line shape of the Z⁰ and determination of electroweak parameters from its hadronic and leptonic decays," Nucl. Phys. B **417**, 3 (1994) [Erratum-ibid. B **426**, 244 (1994)].
- 629. P. Abreu *et al.* [DELPHI Collaboration], "Study of hard scattering processes in multi-hadron production from gamma gamma collisions at LEP," Z. Phys. C **62**, 357 (1994).
- 630. P. Abreu *et al.* [DELPHI Collaboration], "Interference of neutral kaons in the hadronic decays of the Z^0 ," Phys. Lett. B **323**, 242 (1994).
- 631. P. Abreu *et al.* [DELPHI Collaboration], "Invariant mass dependence of particle correlations in hadronic final states from the decay of the Z^0 ," Z. Phys. C **63**, 17 (1994).
- 632. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the $B^0 \bar{B}^0$ mixing using the average electric charge of hadron jets in Z^0 decays," Phys. Lett. B **322**, 459 (1994).
- 633. F. Abe *et al.* [CDF Collaboration], "A Search for the top quark decaying to charged Higgs in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **72**, 1977 (1994).
- 634. P. Abreu *et al.* [DELPHI Collaboration], "Production of Λ and $\Lambda\bar{\Lambda}$ correlations in the hadronic decays of the Z^0 ," Phys. Lett. B **318**, 249 (1993).
- 635. P. Abreu et al. [DELPHI Collaboration], Z. Phys. C **61**, 407 (1994).
- 636. P. Abreu *et al.* [DELPHI Collaboration], "Limits on the production of scalar leptoquarks from Z^0 decays at LEP," Phys. Lett. B **316**, 620 (1993).
- 637. F. Abe *et al.* [CDF Collaboration], "Measurement of the anti-proton proton total cross-section at $\sqrt{s} = 546$ -GeV and 1800-GeV," Phys. Rev. D **50**, 5550 (1994).
- 638. F. Abe *et al.* [CDF Collaboration], "Measurement of small angle anti-proton proton elastic scattering at $\sqrt{s} = 546$ -GeV and 1800-GeV," Phys. Rev. D **50**, 5518 (1994).
- 639. F. Abe et al. [CDF Collaboration], "Measurement of \bar{p} p single diffraction dissociation at $\sqrt{s} = 546$ -GeV and 1800-GeV," Phys. Rev. D **50**, 5535 (1994).
- 640. F. Abe *et al.* [CDF Collaboration], "Study of four jet events and evidence for double parton interactions in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **47**, 4857 (1993).

- 641. F. Abe *et al.* [CDF Collaboration], "Inclusive χ_c and b quark production in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **71**, 2537 (1993).
- 642. P. Abreu *et al.* [DELPHI Collaboration], "Search for Z^0 decays to two leptons and a charged particle anti-particle pair," Nucl. Phys. B **403**, 3 (1993).
- 643. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the mean lifetimes of charged and neutral *B* hadrons," Phys. Lett. B **312**, 253 (1993).
- 644. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of D meson production in Z^0 hadronic decays," Z. Phys. C **59**, 533 (1993) [Erratum-ibid. C **65**, 709 (1993)].
- 645. F. Abe *et al.* [CDF Collaboration], "Measurement of bottom quark production in 1.8-TeV $p \bar{p}$ collisions using semileptonic decay muons," Phys. Rev. Lett. **71**, 2396 (1993).
- 646. P. Abreu *et al.* [DELPHI Collaboration], "Determination of α_s from the scaling violation in the fragmentation functions in e^+ e^- annihilation," Phys. Lett. B **311**, 408 (1993).
- 647. P. Abreu *et al.* [DELPHI Collaboration], "Determination of α_s for *b* quarks at the Z^0 resonance," Phys. Lett. B **307**, 221 (1993).
- 648. F. Abe *et al.* [CDF Collaboration], "Measurement of Drell-Yan electron and muon pair differential cross-sections in \bar{p} p collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **49**, 1 (1994).
- 649. F. Abe *et al.* [CDF Collaboration], "A Search for first generation leptoquarks in \bar{p} p collisions at $\sqrt{s} = 1.8$ TeV," Phys. Rev. D **48**, 3939 (1993).
- 650. F. Abe *et al.* [CDF Collaboration], "Measurement of the bottom quark production cross-section using semileptonic decay electrons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **71**, 500 (1993).
- 651. F. Abe *et al.* [CDF Collaboration], "Measurement of jet multiplicity in W events produced in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **70**, 4042 (1993).
- 652. P. Abreu *et al.* [DELPHI Collaboration], "Determination of α_s using the next-to-leading log approximation of QCD," Z. Phys. C **59**, 21 (1993).
- 653. P. Abreu et al. [DELPHI Collaboration], "Measurement of Λ_b production and lifetime in Z^0 hadronic decays," Phys. Lett. B **311**, 379 (1993).
- 654. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the triple gluon vertex from four-jet events at LEP," Z. Phys. C **59**, 357 (1993).
- 655. F. Abe *et al.* [CDF Collaboration], "The center-of-mass angular distribution of prompt photons produced in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **71**, 679 (1993).
- 656. ** P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the tau lifetime," Phys. Lett. B **302**, 356 (1993).

- 657. F. Abe *et al.* [CDF Collaboration], "Measurement of the dijet mass distribution in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **48**, 998 (1993).
- 658. F. Abe *et al.* [CDF Collaboration], "Measurement of the cross-section for production of two isolated prompt photons in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **70**, 2232 (1993).
- 659. F. Abe *et al.* [CDF Collaboration], "A prompt photon cross-section measurement in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **48**, 2998 (1993).
- 660. P. Abreu *et al.* [DELPHI Collaboration], "A Study of $B^0\bar{B}^0$ mixing using semileptonic decays of B hadrons produced from Z^0 ," Phys. Lett. B **301**, 145 (1993).
- 661. P. Abreu *et al.* [DELPHI Collaboration], "A Search for lepton flavor violation in \mathbb{Z}^0 decays," Phys. Lett. B **298**, 247 (1993).
- 662. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of inclusive production of light meson resonances in hadronic decays of the Z^0 ," Phys. Lett. B **298**, 236 (1993).
- 663. F. Abe *et al.* [CDF Collaboration], "Comparison of jet production in \bar{p} p collisions at \sqrt{s} = 546-GeV and 1800-GeV," Phys. Rev. Lett. **70**, 1376 (1993).
- 664. F. Abe *et al.* [CDF Collaboration], "Search for $\Lambda_b \to J/\psi \Lambda^0$ in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **47**, 2639 (1993).
- 665. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of B meson production and lifetime using D lepton- events in Z^0 decays," Z. Phys. C **57**, 181 (1993).
- 666. ** N. Bingefors *et al.*, "The DELPHI microvertex detector," Nucl. Instrum. Meth. A **328**, 447 (1993).
- 667. F. Abe et al. [CDF Collaboration], "Inclusive J/ψ , $\psi(2S)$ and b quark production in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **69**, 3704 (1992).
- 668. F. Abe *et al.* [CDF Collaboration], "A Measurement of jet shapes in $p \bar{p}$ collisions at \sqrt{s} = 1.8-TeV," Phys. Rev. Lett. **70**, 713 (1993).
- 669. F. Abe *et al.* [CDF Collaboration], "Search for squarks and gluinos from \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **69**, 3439 (1992).
- 670. F. Abe *et al.* [CDF Collaboration], "Limit on the rare decay $W^{\pm} \to \gamma \pi^{\pm}$ in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **69**, 2160 (1992).
- 671. P. Abreu *et al.* [DELPHI Collaboration], "Multiplicity fluctuations in hadronic final states from the decay of the Z^0 ," Nucl. Phys. B **386**, 471 (1992).
- 672. F. Abe *et al.* [CDF Collaboration], "The Dijet angular distribution in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **69**, 2896 (1992).

- 673. P. Abreu *et al.* [DELPHI Collaboration], "Evidence for B_s^0 meson production in Z^0 decays," Phys. Lett. B **289**, 199 (1992).
- 674. F. Abe *et al.* [CDF Collaboration], "Limits on the production of massive stable charged particles," Phys. Rev. D **46**, 1889 (1992).
- 675. P. Abreu *et al.* [DELPHI Collaboration], "Bose-Einstein correlations in the hadronic decays of the Z^0 ," Phys. Lett. B **286**, 201 (1992).
- 676. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the partial width of the Z^0 into $b\bar{b}$ final states using their semileptonic decays," Z. Phys. C **56**, 47 (1992).
- 677. P. Abreu *et al.* [DELPHI Collaboration], "Charged particle multiplicity distributions for fixed number of jets in Z^0 hadronic decays," Z. Phys. C **56**, 63 (1992).
- 678. P. Abreu *et al.* [DELPHI Collaboration], "A Study of the decays of tau leptons produced on the Z resonance at LEP," Z. Phys. C **55**, 555 (1992).
- 679. F. Abe *et al.* [CDF Collaboration], "A Measurement of the production and muonic decay rate of W and Z bosons in p \bar{p} collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **69**, 28 (1992).
- 680. P. Abreu *et al.* [DELPHI Collaboration], "Determination of α_s in second order *QCD* from hadronic *Z* decays," Z. Phys. C **54**, 55 (1992).
- 681. F. Abe *et al.* [CDF Collaboration], "Measurement of the isolated prompt photon cross-sections in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 2734 (1992).
- 682. F. Abe *et al.* [CDF Collaboration], "A Measurement of the *B* meson and *b* quark cross-sections at $\sqrt{s} = 1.8$ -TeV using the exclusive decay $B^{\pm} \to J/\psi K^{\pm}$," Phys. Rev. Lett. **68**, 3403 (1992).
- 683. F. Abe *et al.* [CDF Collaboration], "Measurement of the ratio $\sigma \cdot \mathcal{B}(W \to \tau \nu) / \sigma \cdot \mathcal{B}(W \to e\nu)$, in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 3398 (1992).
- 684. P. Abreu *et al.* [DELPHI Collaboration], "Charged particle multiplicity distributions in restricted rapidity intervals in Z^0 hadronic decays," Z. Phys. C **52**, 271 (1991).
- 685. F. Abe *et al.* [CDF Collaboration], "A Limit on the top quark mass from proton antiproton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **45**, 3921 (1992).
- 686. F. Abe *et al.* [CDF Collaboration], "Lepton asymmetry in W decays from \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 1458 (1992).
- 687. F. Abe *et al.* [CDF Collaboration], "Measurement of $B^0\bar{B}^0$ mixing at the Fermilab Tevatron collider," Phys. Rev. Lett. **67**, 3351 (1991).
- 688. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the Z^0 branching fraction to b quark pairs using the boosted sphericity product," Phys. Lett. B **281**, 383 (1992).

- 689. F. Abe *et al.* [CDF Collaboration], "Measurement of the Z (p_T) distribution in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **67**, 2937 (1991).
- 690. P. Abreu *et al.* [DELPHI Collaboration], "Production of strange particles in the hadronic decays of the Z^0 ," Phys. Lett. B **275**, 231 (1992).
- 691. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the $b\bar{b}$ forward backward asymmetry using the semileptonic decay into muons," Phys. Lett. B **276**, 536 (1992).
- 692. P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of $\sin^2 \theta_W$ from the charge asymmetry of hadronic events at the Z^0 peak," Phys. Lett. B **277**, 371 (1992).
- 693. P. Abreu *et al.* [DELPHI Collaboration], "Multiplicity dependence of mean transverse momentum in e^+e^- annihilations at LEP energies," Phys. Lett. B **276**, 254 (1992).
- 694. P. Abreu *et al.* [DELPHI Collaboration], "Searches for heavy neutrinos from Z decays," Phys. Lett. B **274**, 230 (1992).
- 695. F. Abe *et al.* [CDF Collaboration], "Properties of events with large total transverse energy produced in proton anti-proton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **45**, 2249 (1992).
- 696. F. Abe *et al.* [CDF Collaboration], "A Search for new gauge bosons in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 1463 (1992).
- 697. F. Abe *et al.* [CDF Collaboration], "A Lower limit on the top quark mass from events with two leptons in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 447 (1992).
- 698. P. Abreu *et al.* [DELPHI Collaboration], "Study of final state photons in hadronic Z^0 decay and limits on new phenomena," Z. Phys. C **53**, 555 (1992).
- 699. F. Abe *et al.* [CDF Collaboration], "The Topology of three jet events in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **45**, 1448 (1992).
- 700. P. Abreu *et al.* [DELPHI Collaboration], "Measurement of the average lifetime of *B* hadrons," Z. Phys. C **53**, 567 (1992).
- 701. P. Abreu *et al.* [DELPHI Collaboration], "A Search for neutral Higgs particles in \mathbb{Z}^0 decays," Nucl. Phys. B **373**, 3 (1992).
- 702. P. Abreu *et al.* [DELPHI Collaboration], "Search for scalar leptoquarks from Z^0 decays," Phys. Lett. B **275**, 222 (1992).
- 703. F. Abe *et al.* [CDF Collaboration], "Inclusive jet cross-section in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **68**, 1104 (1992).
- 704. F. Abe et al. [CDF Collaboration], "Search for $W' \to e\nu$ and $W' \to \mu\nu$ in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **67**, 2609 (1991).

- 705. ** P. Abreu *et al.* [DELPHI Collaboration], "A Measurement of the lifetime of the tau lepton," Phys. Lett. B **267**, 422 (1991).
- 706. P. Abreu et al. [DELPHI Collaboration], "The reaction $e^+e^- \to \gamma\gamma(\gamma)$ at Z^0 energies," Phys. Lett. B **268**, 296 (1991).
- 707. P. Abreu *et al.* [DELPHI Collaboration], "Study of orientation of three jet events in Z^0 hadronic decays using the DELPHI detector," Phys. Lett. B **274**, 498 (1992).
- 708. F. Abe *et al.* [CDF Collaboration], "Measurement of the e^+e^- invariant mass distribution in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **67**, 2418 (1991).
- 709. F. Abe *et al.* [CDF Collaboration], "A Determination of $\sin^2 \theta$ from the forward backward asymmetry in $p\bar{p} \to Z^0 X \to e^+ e^- X$ interactions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **67**, 1502 (1991).
- 710. P. Abreu *et al.* [DELPHI Collaboration], "Search for excited charged leptons in \mathbb{Z}^0 decays," Z. Phys. C **53**, 41 (1992).
- 711. P. Abreu *et al.* [DELPHI Collaboration], "Determination of Z^0 resonance parameters and couplings from its hadronic and leptonic decays," Nucl. Phys. B **367**, 511 (1991).
- 712. F. Abe *et al.* [CDF Collaboration], "Measurement of QCD jet broadening in $p \bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **44**, 601 (1991).
- 713. P. Abreu *et al.* [DELPHI Collaboration], "A Study of the reaction $e^+e^- \to \mu^+\mu^-$ around the Z^0 pole," Phys. Lett. B **260**, 240 (1991).
- 714. F. Abe *et al.* [CDF Collaboration], "Measurement of the Wp_T distribution in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **66**, 2951 (1991).
- 715. P. Abreu *et al.* [DELPHI Collaboration], "Search for low mass Higgs bosons produced in Z^0 decays," Z. Phys. C **51**, 25 (1991).
- 716. F. Abe et al. [CDF Collaboration], "A measurement of $\sigma \cdot \mathcal{B}(W \to e\nu)$ and $\sigma \cdot \mathcal{B}(Z^0 \to e^+e^-)$ in \bar{p} p collisions at $\sqrt{s} = 1800$ -GeV," Phys. Rev. D 44, 29 (1991).
- 717. P. Abreu *et al.* [DELPHI Collaboration], "Experimental study of the triple gluon vertex," Phys. Lett. B **255**, 466 (1991).
- 718. ** F. Abe *et al.* [CDF Collaboration], "A measurement of the W boson mass in 1.8-TeV \bar{p} p collisions," Phys. Rev. D **43**, 2070 (1991).
- 719. ** F. Abe *et al.* [CDF Collaboration], "A measurement of the W boson mass," Phys. Rev. Lett. **65**, 2243 (1990).
- 720. F. Abe *et al.* [CDF Collaboration], "Top quark search in the electron + jet channel in proton anti-proton collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. D **43**, 664 (1991).

- 721. F. Abe *et al.* [CDF Collaboration], "Search for new heavy quarks in electron muon events at the Fermilab Tevatron collider," Phys. Rev. Lett. **64**, 147 (1990).
- 722. F. Abe *et al.* [CDF Collaboration], "Search for a light Higgs boson at the Tevatron proton anti-proton collider," Phys. Rev. D **41**, 1717 (1990).
- 723. F. Abe et al. [CDF Collaboration], "Measurement of the ratio $\sigma(W \to e\nu) / \sigma(Z \to ee)$ in \bar{p} p collisions at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **64**, 152 (1990).
- 724. F. Abe *et al.* [CDF Collaboration], "A search for the Top quark in the reaction $\bar{p}p \rightarrow$ electron + jets at $\sqrt{s} = 1.8$ -TeV," Phys. Rev. Lett. **64**, 142 (1990).
- 725. ** F. Abe *et al.* [CDF Collaboration], "Measurement of the mass and width of the Z^0 boson at the Fermilab Tevatron," Phys. Rev. Lett. **63**, 720 (1989).
- 726. F. Abe *et al.* [CDF Collaboration], "A measurement of W boson production in 1.8-TeV \bar{p} p collisions," Phys. Rev. Lett. **62**, 1005 (1989).
- 727. F. Abe *et al.* [CDF Collaboration], "The CDF detector: An overview," Nucl. Instrum. Meth. A **271**, 387 (1988).

Conference proceedings

- 1. M. Mikuz, D. Asner, M. Barbero, V. Bellini, V. Belyaev, J. M. Brom, M. Bruzzi, D. Chren et al., "Diamond detectors," Proceedings of **VERTEX2010**, 024 (2010).
- 2. ** I. Dolenc *et al.*, "ATLAS diamond Beam Conditions Monitor," PoS **VERTEX2008**, 012 (2008).
- 3. ** W. Trischuk [RD42 Collaboration], "Recent Advances in Diamond Detectors," Published in Proceedings of the 34th International Conference on High-Energy Physics (ICHEP 08), Philadelphia, Aug 2008, arXiv:0810.3429.
- 4. ** C. P. Hays, B. Jayatilaka, A. Kotwal, L. Nodulman, O. Stelzer-Chilton, W. Trischuk and I. Vollrath, "First measurement of the W boson mass with CDF in run II," *Published in the proceedings of the International Europhysics Conference on High Energy Physics (EPS-HEP2007), Manchester, England, July 2007*, J. Phys. Conf. Ser. **110**, 122019 (2008).
- 5. ** W. Trischuk [CDF-II Collaboration], "Tevatron Results on W Boson Mass and Width," Nuclear Physics **B** (Proc. Suppl.) 177-178, 22 (2008). Published in Proceedings of the Hadron Collider Physics Symposium (HCP 2007), Elba, Italy, 20-26 May 2007.
- 6. ** W. Trischuk [RD42 Diamond Detector Collaboration], "Recent Developments In Diamond Particle Detectors," Published in Proceedings of the 32nd International Conference on High-Energy Physics (ICHEP 04), Beijing, China, 16-22 Aug 2004.

- 7. ** W. Trischuk [BELLE Collaboration], "Measurement of *CP* Asymmetries at Belle" arXiv:hep-ex/0212059. *Published in the proceedings of the 2002 SLAC Summer Institute, Topical Conference.*
- 8. ** W. Trischuk [CDF Collaboration], "Experimental study of sin(2β) and sin(2α)," FERMILAB-CONF-00-044-E Published in the proceedings of 3rd International Conference on B Physics and CP Violation (BCONF99), Taipei, Taiwan, 3-7 Dec 1999.
- 9. ** W. Trischuk [CDF collaboration], "Rare B decays from the CDF experiment," Nucl. Phys. Proc. Suppl. **75B**, 89 (1999).
- 10. ** W. Trischuk, "Semiconductor trackers for future particle physics detectors," Nucl. Instrum. Meth. A **419**, 251 (1998).
- 11. ** W. Trischuk [RD42 Collaboration], "Diamond detectors for the LHC," Prepared for International Europhysics Conference on High-energy Physics (HEP 95), Brussels, Belgium, 27 Jul 2 Aug 1995.
- 12. ** W. Trischuk [DELPHI Collaboration], "A review of the tau lifetime," CERN-PPE-92-197 Invited talk at Workshop on Tau Lepton Physics, Columbus, OH, Sep 8-11, 1992.
- 13. ** W. Trischuk [DELPHI Collaboration], "B and D lifetimes at LEP," CERN-PPE-91-128 Invited Talk at the 4th Int. Symp. on Heavy Flavor Physics, Orsay, France, Jun 25-28, 1991.
- 14. ** W. Trischuk [CDF Collaboration], "Preliminary electroweak results from CDF electron data," Nucl. Phys. Proc. Suppl. **16**, 205 (1990).

Non-Refereed publications

- 1. M. Butler *et al.*, "The Subatomic Universe: Canada in the Age of Discovery 2011-2016", Report of the NSERC Long-Range Planning Committee, March 2012.
- 2. ** K.J. Ragan *et al.*, "Perspectives on Subatomic Physics in Canada 2006-2016", Report of the NSERC Long-Range Planning Committee, December 2006.
- 3. W. Adam *et al.* [RD42 Collaboration], "Development of diamond tracking detectors for high luminosity experiments at the LHC," CERN-LHCC-2002-010.
- 4. ** K. Anikeev *et al.*, "B physics at the Tevatron: Run II and beyond," hep-ph/0201071, proceedings of FNAL Run II B Physics workshop.
- 5. ** W. Adam *et al.* [RD42 Collaboration], "Development of diamond tracking detectors for high luminosity experiments at the LHC," CERN-LHCC-2000-011.
- 6. R. Blair *et al.* [CDF-II Collaboration], "The CDF-II detector: Technical design report," FERMILAB-PUB-96-390-E.

- 7. ** M. H. Nazare *et al.* [RD42 Collaboration], "R & D proposal: Development of diamond tracking detectors for high luminosity experiments at the LHC," CERN-DRDC-94-21.
- 8. ** W. Trischuk, "A measurement of the W boson mass in 1.8-TeV proton anti proton collisions," UMI-90-35576, PhD Thesis, Harvard University (1990).