




## FLAVOUR MIXING BY WEAK FORCE

WEAK FORCE DOES NOT CONSERVE QUARK FLAVOUR  
JUST FLIPS LEPTONS WITHIN DOUBLETS



CHANGES FLAVOUR OF QUARKS



THINK ABOUT FIRST 2 GENERATIONS

$$\begin{pmatrix} u \\ d \end{pmatrix} \quad \begin{pmatrix} c \\ s \end{pmatrix}$$

- STATES OF DEFINITE
- MASS
  - COLOUR
  - FLAVOUR

→ EIGENSTATES OF MASS

$$M|u\rangle = m_u|u\rangle$$

→ EIGENSTATES OF COLOUR FORCE

$$C|u\rangle = c_u|u\rangle$$

THESE CANNOT BE THE EIGENSTATES THAT WEAK INTERACTION SEES → IT DOES NOT CONSERVE QUARK FLAVOUR

EIGENSTATES OF WEAK INTERACTION ARE  
A MIXTURE OF DEFINITE FLAVOUR STATES

→ THIS IS WHY WEAK INTERACTION CAN INDUCE  
TRANSITIONS BETWEEN STATES OF DEFINITE FLAVOUR

$$\begin{pmatrix} u \\ d \end{pmatrix}$$

↑  
COLOUR EIGENSTATE  
= MASS EIGENSTATE

$$\begin{pmatrix} u \\ d \cos \theta_c + s \sin \theta_c \end{pmatrix}$$

WEAK EIGENSTATES

$$\begin{array}{l} d \rightarrow u \quad \alpha \quad \cos^2 \theta_c \\ s \rightarrow u \quad \alpha \quad \sin^2 \theta_c \end{array}$$


$$\begin{pmatrix} d \\ s \end{pmatrix}_{\text{WEAK}} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} d \\ s \end{pmatrix}_{\text{COLOUR}}$$

↑  
CABIBBO ANGLE

MAINLY WEAK INTERACTION INDUCES



$d$  &  $s$  BEING MIXED — CAN INDUCE




MODIFIES UNIVERSAL FERMION WEAK COUPLING

→ MEASUREMENTS CONSISTENT WITH  
UNIVERSAL CABBIBO ANGLE

$\theta_c$

$$\mu^- \rightarrow \nu_\mu e^- \bar{\nu}_e$$




STRENGTH

$$1 \times G_F^2$$

$$d \rightarrow u e^- \bar{\nu}_e$$

( $n \rightarrow p e^- \nu$ )




$$\sim 0.95$$

( $\cos^2 \theta_c$ )

$$s \rightarrow u e^- \bar{\nu}_e$$

( $\Lambda \rightarrow p e^- \nu$ )  
( $Br \sim 10^{-3}$ )




$$\sim 0.05$$

( $\sin^2 \theta_c$ )

$$s \rightarrow u d \bar{u}$$



( $\Lambda \rightarrow p \pi^-$ )



$$\sim 0.09$$

( $\sin \theta_c \cos \theta_c$ )

Z<sup>0</sup> DOES NOT INDUCE DECAYS - WHY?




PARITY VIOLATION

AMP  $\sim \frac{G}{\sqrt{2}} \cdot \sin \theta_c \cdot f_K m_\mu \bar{\nu} \delta_S \mu$

DECAY  $\Gamma \sim \frac{G^2}{8\pi} \sin^2 \theta_c f_K^2 m_K m_\mu^2 (1 - m_\mu^2/m_K^2)$

AGREES WITH EXPERIMENT  $\tau \sim 10^{-8} s, BR \sim 64\%$




SHOULD BE ABOUT SAME  $\Gamma$  AS THE W<sup>-</sup> DIAGRAM


EXPERIMENTALLY  $\tau_{K_L} = 5 \times 10^{-8} s$   
 $BR = 9.5 \times 10^{-9}$

NO FLAVOUR CHANGING NEUTRAL CURRENTS

# NO FIRST ORDER FLAVOUR CHANGING NEUTRAL CURRENT



BUT  $W^\pm$  CAN INDUCE AT HIGHER ORDER





$$\frac{\Gamma(K_L^0 \rightarrow \mu\mu)}{\Gamma(K^+ \rightarrow \mu\nu)} \approx \left( \frac{3\sqrt{2}\alpha}{\pi} \right)^2 \rightarrow BR(K_L^0 \rightarrow \mu^+\mu^-) \approx 3 \times 10^{-4}$$

cf EXPERIMENT  $BR \sim 10^{-9}$  } SOME SUPPRESSION MECHANISM IS AT WORK HERE


BUT NEUTRAL CURRENTS DO EXIST.

THE FOLLOWING  $\nu$  INTERACTIONS OBSERVED WITH EXPECTED WEAK COUPLING STRENGTH



NOTICE THAT THESE INTERACTIONS DO NOT CHANGE QUARK FLAVOUR FROM INITIAL TO FINAL STATE

WHY DOES  $Z^0$  NOT INDUCE






FROM


$$d \cos\theta + s \sin\theta$$


WE CAN WRITE THE TRANSITION AMPLITUDES


$$\sim u \bar{d} \cos\theta$$





$$u \bar{s} \sin\theta$$



BUT ALSO HAVE


$$d \cos\theta \quad s \sin\theta$$




$$\begin{pmatrix} u \\ d \cos \theta & s \sin \theta \end{pmatrix}$$


EXPAND OUT ALL POSSIBLE TRANSITIONS FOR  $Z^0$

$$u\bar{u} + d\bar{d} \cos^2 \theta + s\bar{s} \sin^2 \theta$$



FLAVOUR CONSERVING  
OBSERVED IN  $V$  SCATTERING

$$s\bar{d} \sin \theta \cos \theta + \bar{d}s \sin \theta \cos \theta$$




FLAVOUR CHANGING  
NOT OBSERVED IN DECAY


ABSENCE OF  $Z^0$  DECAYS LED

GLASHOW, ILIO POULOUS & MAIANI TO MAKE THE  
FOLLOWING PREDICTION BEFORE C-QUARK DISCOVERY

$$\begin{pmatrix} u \\ d \end{pmatrix} \quad \begin{pmatrix} c \\ s \end{pmatrix} \quad \rightarrow \quad \text{COLOUR EIGENSTATES}$$

$$\begin{pmatrix} u \\ d \cos \theta + s \sin \theta \end{pmatrix} \quad \begin{pmatrix} c \\ s \cos \theta - d \sin \theta \end{pmatrix}$$


AS THE WEAK EIGENSTATES




$$\left( \begin{array}{c} c \\ s \cos \theta - d \sin \theta \end{array} \right)$$

$$\left( \begin{array}{c} \bar{c} \\ \bar{s} \cos \theta - \bar{d} \sin \theta \end{array} \right)$$

NFC = NO FLAVOUR CHANGE

FC = FLAVOUR CHANGE




WRITE OUT TRANSITION AMPLITUDES

$$u\bar{u} + c\bar{c} + (\bar{d}d + s\bar{s})\cos^2\theta + (s\bar{s} + d\bar{d})\sin^2\theta$$

$$+ (s\bar{d} + \bar{s}d - \bar{s}d - s\bar{d})\cos\theta\sin\theta$$



→ CONSERVES FLAVOUR  
→ FLAVOUR CHANGING PART VANISHES




THESE CANNOT  
MEDIATE DECAYS

$$m_u = m_{\bar{u}}$$

$$m_c = m_{\bar{c}}$$




CANCELLATION



# GIM MECHANISM PREDICTED $m_{\text{CHARM}}$

FOR  $K_L^0 \rightarrow \mu^+ \mu^-$  NOW HAVE TWO 2ND ORDER DIAGRAMS



$$BR(K_L \rightarrow \mu\mu) \sim 7 \times 10^{-5} \frac{m_c^2 - m_u^2}{M_W^2} \ln \frac{M_W^2}{m_u^2}$$

PREDICTED  $\rightarrow m_c \approx 1.5 \frac{\text{GeV}}{c^2}$  ✓

GENERALLY TRUE THAT THESE BOX DIAGRAMS  
ARE DOMINATED BY HEAVIEST QUARK THAT  
CAN CONTRIBUTE TO THE INTERNAL LOOP


$m_c \rightarrow K$  DECAYS

$m_t \rightarrow B^0 \bar{B}^0$  MIXING


$m_H \rightarrow$  RADIATIVE CORRECTIONS

RARE DECAYS CAN ACCESS HIGHER MASS  
SCALES THAN DIRECT PRODUCTION AT  
ACCELERATORS

$D^0 \bar{D}^0$



$B_d^0 \bar{B}_d^0$



$m_t > 150 \frac{\text{GeV}}{c^2}$

$e^+e^- \rightarrow b\bar{b}$   
@ 10 GeV

$B_s^0 \bar{B}_s^0$

