

LECTURE 9: Wave Packets

Goal of the lecture: describe the wave function of a quantum particle using wave packets

What I expect you to learn:

- what are waves packets
- what is the group and phase velocities of wave packets
- what are Fourier Transforms
- how to express the wave function in momentum space
- how to calculate Fourier Transforms

(This roughly corresponds to section 2.4, appendix A of textbook)

A quick recap

(2)

→ We related wave-type quantities to a particle's energy and momentum:

$$E = \hbar \omega$$
$$p = \hbar k$$

→ For a free particle we have: $E = \frac{p^2}{2m} = \frac{mV^2}{2}$

which means we the following relation between ω and k :

$$\omega = \frac{\hbar k^2}{2m}$$

→ we found an equation that obeyed this relation:

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V\psi = i\hbar \frac{\partial \psi}{\partial t}$$

A simple solution to this equation is

$$\psi(x,t) = A e^{i(kx - \omega t)}$$

A quick recap (cont.)



(3)

- we saw that in order to account for interference effects, we must be able to superimpose wave functions e.g.

$$\psi = c_1 \psi_1 + c_2 \psi_2 + \dots \text{(superposition principle)}$$



- As a first step on our way to superimposing wave functions we considered the case of a string attached at two ends:




A quick RECAP (CONT.)

(4)

- WE SAW THAT WE COULD EXPRESS AN ARBITRARY WAVE FORM ON THE STRING AS A SUM OF SINE AND COSINE WAVES i.e. A FOURIER SERIES:

$$\textcircled{1} f(x) = \frac{A_0}{2} + \sum_{n=1}^{\infty} A_n \sin nx + \sum_{n=1}^{\infty} B_n \cos nx$$

FOR EXAMPLE WE SAW :



→ we found the amplitudes for this example

note that $\textcircled{1}$ has discrete values of k

We need to work with a continuum of k values

Wave Functions for Fixed p

(5)

For a fixed momentum, we have

$$\begin{aligned}\psi(x,t) &= A e^{i(kx - \omega t)} \\ &= A e^{i/k(px - Et)}\end{aligned}$$

Note that

$$\begin{aligned}-i\hbar \frac{\partial}{\partial x} \psi &= p\psi & i\hbar \frac{\partial}{\partial t} \psi &= E\psi\end{aligned}$$

↑ differential operators ↓

Probability interpretation of $|\psi|^2 \Rightarrow$

$$\int_{-\infty}^{\infty} |\psi(x,t)|^2 dx = 1$$

The plane wave cannot be normalized to unity

WAVE PACKETS IN 1 DIMENSION

(6)

WE NEED TO SUM PLANE WAVES... THE MOST GENERAL EXPRESSION IS:

$$\psi(x, t) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \varphi(p) e^{i/\hbar}(px - Et) dp$$

→ Amplitude for wave of momentum p (or $\hbar k$) → can be complex

Assume that $\varphi(p)$ is peaked about $p = p_0$ i.e. falls to 0 outside $[p_0 - \Delta p, p_0 + \Delta p]$

Take $B(p) = px - Et$

$|B(x, t)|$ will be largest when $B(p)$ is almost constant around $p = p_0$

→ remember $e^{i\theta} = \cos\theta + i\sin\theta$

⇒ get max of $|B(x, t)|$ when $\frac{dB}{dp} \approx 0$

WAVE PACKETS (CONT.)

(9)

→ The max of $|2t(x,t)|$ will correspond to the center of the wave packet

$$\text{We have } E = \frac{p^2}{2m}, \quad B = px - Et$$

$$\frac{dB}{dp} = x - \frac{p}{m} t = x - vt = 0$$

$x = vt \quad \hookrightarrow v_g$

v_g : velocity of the group of plane waves

$$v_g = \frac{dw}{dk} = \frac{dE}{dp}, \quad v_p = \text{velocity of plane waves}$$

$$A e^{i(kx - \omega t)} = A e^{iK(x - \frac{\omega}{K}t)} \Rightarrow v_p = \frac{\omega}{K}$$

Let's take a look here:

<http://webphysics.davidson.edu/Applets/superposition/GroupVelocity.ht>

WAVE PACKETS (CONT.)

(8)

WE CAN EXPAND $E(p)$ ABOUT $p_0 \rightarrow$ TAYLOR EXPANSION


$$E(p) = p^2/2m$$

$$E(p) = E(p_0) + (p-p_0) \left. \frac{dE(p)}{dp} \right|_{p=p_0} + \frac{1}{2} (p-p_0)^2 \left. \frac{d^2E(p)}{dp^2} \right|_{p=p_0} + \dots$$

$$= \frac{p_0^2}{2m} + (p-p_0) \frac{p_0}{m} + \frac{(p-p_0)^2}{2m} + \dots$$

$$\psi(x,t) = \underbrace{e^{i\frac{1}{\hbar}(p_0 x - E(p_0)t)}}_{\text{plane wave}} \cdot \underbrace{\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{i(\frac{p-p_0}{\hbar}(x - v_g t) + \frac{1}{2\hbar}(p-p_0)^2 t)} \phi(p) dp}_{\text{envelope function}}$$


\hookrightarrow neglect: $\Delta p^2 t \ll 1$



FOURIER TRANSFORM INTERLUDE

(4)

$$f(x) = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n \sin \frac{n\pi x}{L} + B_n \cos \frac{n\pi x}{L}$$



we can also write:

$$f(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} C_n e^{in\pi x/L}$$

For non-periodic functions we take $L \rightarrow \infty$

as $L \rightarrow \infty$, differences between terms in the series become smaller:

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} C_n e^{in\pi x/L} dn$$

let $k = n\pi/L$ and $g(k) = L \frac{C_n}{\pi}$, $dk = dn \cdot \frac{\pi}{L}$

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(k) e^{ikx} dk, \quad g(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx$$

Fourier Transforms (cont.)

(10)

Physically what does this all mean?

For $T=0$, we have

$$\psi(x) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \varphi(p) e^{ipx/\hbar} dp$$

→ probability amplitude for finding particle
→ sum of plane waves with amplitude $\varphi(p)$



$$\varphi(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \psi(x) e^{-ipx/\hbar} dx$$

→ probability amplitude for finding particle with momentum p
→ sum of plane waves with amplitude $\psi(x)$
→ momentum space wave function

Fourier Transforms (cont.)

(11)

$$a) \quad T=0$$



For a particle whose prob. amplitude dist. looks like this, what are the amplitudes of its components? plane waves?

At $T=0$, the particle is most probably at x_0 and it's velocity is given by the group velocity:

$$v_g = \left. \frac{dE}{dp} \right|_{x=x_0}$$

