



CARTOON OF MODERN ACCELERATOR COMPLEX


Large Hadron Collider at CERN

Circumference 26.7 km (16.6 miles)



BUILDING THE LHC





ACCELERATORS


- GEIGER & MARSDEN - RADIOACTIVE SOURCE
7 MeV α
- SIZE & SHAPE OF NUCLEUS 40 MeV p
- SIZE & SHAPE OF PROTON 500 MeV e^-
- PATTERN OF π , p, Δ^{++} , K^+ MASSES 10 GeV p
- DISCOVERY OF QUARKS 50 GeV e^-
- DISCOVERY OF TOP QUARK 1000 GeV p
- DISCOVERY OF HIGGS BOSON 14000 GeV p
- AS WE WANT TO RESOLVE SMALLER STRUCTURES
PRODUCE HIGHER MASS
- INCREASE ACCELERATOR ENERGY

TWO PURPOSES

- PRODUCE MASSIVE UNSTABLE PARTICLES
CONVERT KINETIC ENERGY OF BEAMS \rightarrow MASS

$$E = mc^2 \rightarrow m = E/c^2$$


- PROBE SMALLER DISTANCES \rightarrow SCATTERING



DEFINITE MOMENTUM
— PLANE WAVE

$$\Delta p \Delta x \sim \hbar \rightarrow \Delta x \sim \frac{\hbar}{\Delta p}$$

TARGET



$$\Psi = A e^{-i\vec{p} \cdot \vec{r}/\hbar}$$

WAVELENGTH OF BEAM PARTICLES


de BROGLIE $\lambda = h/p$

RESOLVING LENGTH \uparrow BEAM MOMENTUM

SPATIAL RESOLUTION & MOMENTUM TRANSFER

RESOLUTION IN OPTICS

RESOLUTION DEFINED BY




$$\Delta r \sim \frac{\lambda}{\sin \theta}$$


- WAVELENGTH OF LIGHT
- APERTURE

PARTICLE SCATTERING → RESOLVE SMALL OBJECTS USING MATTER WAVES

3-MOMENTUM
- SIMPLICITY



$$\Delta r \approx \frac{\lambda}{\sin \theta} = \frac{h}{p \sin \theta}$$



$$|q| = 2|p| \sin \frac{\theta}{2} \sim p \sin \theta$$

$\Delta r \sim h/q$
 HIGH SPATIAL RESOLUTION ← HIGH MOMENTUM TRANSFER

- PROTON DIAMETER $\sim 1 \text{ fm} = 10^{-15} \text{ m}$
- WHAT MOMENTUM TRANSFER DO WE NEED TO RESOLVE PROTONS INSIDE NUCLEUS?

$$\text{NEED } 1 \text{ fm} = \frac{h}{q} = \frac{hc}{qc} \rightarrow = 1.23 \text{ GeV} \cdot \text{fm}$$

$$qc = 1.23 \text{ GeV} \cdot \text{fm} / 1 \text{ fm}$$

$$q \sim 1 \text{ GeV}/c \quad \text{MOMENTUM TRANSFER}$$

MOMENTUM OF BEAM MUST BE $>$ MOMENTUM TRANSFER

- PRESENT LIMIT ON QUARK DIAMETER $< 10^{-18} \text{ m}$

MOMENTUM TRANSFER


$$\sim 1000 \text{ GeV}/c = 1 \text{ TeV}/c$$

THIS NEEDS AN ACCELERATING VOLTAGE $\sim 10^{12}$ VOLTS

HOW?



LABORATORY & CENTRE OF MASS (MOMENTUM) FRAMES

IN ORDER TO UNDERSTAND HOW TO PRODUCE HIGH MASS OBJECTS \rightarrow SPECIAL RELATIVITY



AS P_{BEAM} INCREASES MORE & MORE ENERGY GOES INTO INCREASING VELOCITY OF CENTRE-OF-MASS AS SEEN IN THE LAB FRAME.

LAB



OBVIOUS FROM
SPECIAL RELATIVITY




- MOTION OF CENTRE-OF-MASS IN LAB IS IRRELEVANT TO PHYSICS
 - ONLY ENERGY IN COFM IS USEFUL FOR NEW PARTICLE PRODUCTION
- DO A LORENTZ BOOST INTO
CM FRAME

"CENTRE-OF-MOMENTUM"



BEFORE

TOTAL MOMENTUM = 0



AFTER

TOTAL MOMENTUM = 0



CONTINUE COMPARISON OF LAB & CMS


LAB: CONSIDER $\pi p \rightarrow \pi N^*$ ← EXCITED PROTON

$$m_{N^*} > m_p > m_\pi$$

IN CMS, BY DEFINITION, π & p
COLLIDE WITH EQUAL & OPPOSITE MOMENTA


CMS:

BEFORE COLLISION



$$\vec{p}_\pi^{CM} = -\vec{p}_p^{CM}$$

AFTER COLLISION



π & N^* AT REST

ALL ENERGY OF RELATIVE
MOTION OF INITIAL π, p
HAS GONE INTO PRODUCING
MASS OF N^*

$$\vec{p}_\pi^{CM} = -\vec{p}_{N^*}^{CM} = 0$$

IN FINAL STATE WHERE π & N^* BOTH AT REST

TOTAL CMS ENERGY

AFTER COLLISION $\left\{ \begin{aligned} W^{CM} &= (m_{\pi} + m_{N^*})c^2 \approx m_{N^*}c^2 \end{aligned} \right.$

NOTICE $(W^{CM})^2 = [m]^2$ $m_{N^*} \gg m_{\pi}$

RELATIVISTIC INVARIANT

TOTAL RELATIVISTIC ENERGY IS CONSERVED IN COLLISION SO

BEFORE COLLISION $\left\{ \begin{aligned} W^{CM} &= E_{\pi}^{CM} + E_p^{CM} = (m_{\pi} + m_{N^*})c^2 \end{aligned} \right.$

OR $(W^{CM})^2 = (E_{\pi}^{CM} + E_p^{CM})^2 \approx m_{N^*}^2 c^2$

RELATIVISTIC INVARIANT

MUST HAVE SAME VALUE IN LABORATORY FRAME.

- REMEMBER ABOUT LORENTZ INVARIANT
(SOME TIMES CALL IT RELATIVISTIC INVARIANT)

- CONSIDER REST MASS m

$$E^2 = p^2 c^2 + m^2 c^4 \rightarrow m^2 c^4 = E^2 - p^2 c^2$$

↑
SAME IN ALL FRAMES

IF WE HAVE A GROUP OF i PARTICLES
4-VECTOR OF EACH IS (E_i, \vec{p}_i)

TOTAL 4-VECTOR:

$$\left(\sum_i E_i\right)^2 - \left(\sum \vec{p}_i\right)^2 c^2 = M^2 c^2$$

↑
ENERGIES

↑
MOMENTUM

↑
LORENTZ
INVARIANT

↑
INVARIANT MASS

INVARIANT MASS OF A GROUP PARTICLES IS A
LORENTZ INVARIANT → JUST LIKE A SINGLE PARTICLE

$$\left(\sum E_i\right)^2 - \left(\sum \vec{p}_i\right)^2 c^2 = M^2 c^4$$

- INVARIANT MASS OF AN \dot{U} PARTICLE SYSTEM IS A LORENTZ INVARIANT YOU CAN CALCULATE IT IN ANY FRAME YOU LIKE
- INVARIANT MASS OF AN \dot{U} PARTICLE SYSTEM IS ONLY EQUAL TO SUM OF MASSES IF RELATIVE MOTION IS ZERO
- THIS MAKES PERFECT SENSE IN SPECIAL RELATIVITY

THE KINETIC ENERGY
DUE TO RELATIVE
MOTION

CONTRIBUTES TO
INVARIANT MASS
(EFFECTIVE MASS)

CONNECTING CM + LAB FRAMES

- USE LORENTZ TRANSFORM? UMM — OK — COMPLEX
- EXPLOIT LORENT INVARIANCE OF INVARIANT MASS

$$M_{LAB}^2 c^4 = M_{CM}^2 c^4$$

CENTRE OF MASS

$$(E_a^{CM} + E_b^{CM})^2 - (\vec{p}_a^{CM} + \vec{p}_b^{CM})^2 c^2 = \left(\begin{array}{c} \text{LAB} \\ E_a + E_b \\ \hline E_{beam} \quad m_{TARG} c^2 \\ \hline p_{beam} \end{array} \right)^2 - (\vec{p}_a^{LAB} + \vec{p}_b^{LAB})^2 c^2 \rightarrow 0$$

$\vec{p}_a^{CM} = -\vec{p}_b^{CM}$

USE $m^2 = E^2 - p^2$

$$W^2 = (E_a^{CM} + E_b^{CM})^2 = 2 E_{beam} m_{TARG} c^2 + (m_{beam}^2 + m_{TARG}^2) c^4$$

COMMON NOTATION

$$W^2 = (E_{TOT}^{CM})^2 = (\sqrt{s})^2$$

TYPICAL

$$W^2 = 2 E_{\text{bm}} \cdot m_{\text{tg}} c^4 + (m_{\text{bm}}^2 + m_{\text{tg}}^2) c^4$$

1000 GeV

1.6 GeV/c²

PROTON-PROTON
FIXED TARGET COLLISIONS

FOR SUCH HIGH ENERGY BEAM - APPROXIMATE


$$W \sim (2 E_{\text{bm}} m_{\text{tg}} c^2)^{1/2}$$

so

$$W \propto \sqrt{E_{\text{beam}}}$$

FIXED TARGET v COLSIDING BEAMS

WILL DISCUSS DETAILS OF ACCELERATORS LATER



- MOST OF p_A GOES INTO MOTION OF CM. IN LAB FRAME
- ACCELERATOR IS IN LAB FRAME

- COLLIDING BEAMS ACCELERATOR IS IN CM FRAME
- AWH OF p_a GOES INTO E_{CM} .

USE 4-MOMENTUM & $C=1$

- 2 PARTICLE COLLISION IN AN ARBITRARY FRAME



$$m_A (E_A, \vec{p}_A) \quad m_B (E_B, \vec{p}_B)$$

- TOTAL (4-MOMENTUM)² p^2 OF SYSTEM

$$p^2 = (p_A + p_B)^2 \quad \leftarrow \text{4-VECTORS}$$

$$p^2 = p_A^2 + p_B^2 + 2p_A p_B = m_A^2 + m_B^2 + 2E_A E_B - 2\vec{p}_A \cdot \vec{p}_B$$

$$p_A^2 = E_A^2 - \vec{p}_A^2 = m_A^2$$

3 VECTOR

$$p^2 = m_A^2 + m_B^2 + 2E_A E_B - 2|\vec{p}_A||\vec{p}_B|\cos\theta$$

SCATTERING ANGLE

$$\text{IN CMS FRAME } \sum \vec{p} = 0$$

$$p^{*2} = E^{*2} - |\vec{p}^*|^2 = E^{*2} \quad \leftarrow \text{(4-VECTOR)}^2$$

* = CMS

→ INVARIANT

$$p^{*2} = E^{*2} = m_A^2 + m_B^2 + 2E_A E_B - 2|\vec{p}_A| |\vec{p}_B| \cos \theta$$


THIS IS TRUE IN ANY FRAME \therefore CAN CHOOSE WHICH FRAME WE EVALUATE IT IN

CHOOSE LAB FRAME $|\vec{p}_B| = 0$; $E_B = m_B$
 STATIONARY TARGET

$$E^{*2} = m_A^2 + m_B^2 + 2m_B E_A \rightarrow E^* \propto \sqrt{E_A} \leftarrow \text{ACCELERATOR ENERGY}$$

• COLLIDING BEAMS: $|\vec{p}_A| = |\vec{p}_B|$ $\cos \theta = -1$

$$E^{*2} = m_A^2 + m_B^2 + 2E_A E_B + 2|\vec{p}_A| |\vec{p}_B|$$



FOR $E \gg m$

$$E^{*2} = 4E_A^2 \quad \text{ACCELERATOR ENERGY}$$

$$E^* = 2E_A \leftarrow \text{ENERGY}$$

$$2E_A^2$$

$$E_A \propto \$$$


$$E_{\text{FIXED}}^* \propto \sqrt{\$}$$

$$E_{\text{COLLIDER}}^* \propto \$$$

ANOTHER USE OF INVARIANT MASS

PARTICLES DECAYING THRU STRONG FORCE $\tau \sim 10^{-23}$ S

HOW CAN ONE MEASURE SUCH A SHORT LIFETIME?



COLLIDE $p \pi^-$

$\pi^- p \rightarrow n \pi^+ \pi^-$


STRONG INTERACTION


CAN PRODUCE VERY SHORT
LIVED INTERMEDIATE STATE

"RESONANCE"


$\tau_{\rho^0} \sim 10^{-23}$ S

TRAVELS $\sim 1f = 10^{-15}$ m





NO INTERMEDIATE STATE
 ENERGY + MOMENTUM SHARED
 BETWEEN $\pi^+ \pi^- \pi$ IN RANDOM
 STATISTICAL FASHIONS
 "PHASE SPACE"



IF INTERMEDIATE STATE





$$E_p = E_{\pi^+} + E_{\pi^-}$$

$$\vec{p}_p = \vec{p}_{\pi^+} + \vec{p}_{\pi^-}$$

$$m_p^2 = [E_p^2 - \vec{p}_p^2]$$

$$= [(E_{\pi^+} + E_{\pi^-})^2 - (\vec{p}_{\pi^+} + \vec{p}_{\pi^-})^2]$$

MASS OF p^0 = INVARIANT MASS OF DAUGHTER
 \bar{u} PAIR



EVIDENCE FOR A $T=0$ THREE-PION RESONANCE*

B. C. Maglić, L. W. Alvarez, A. H. Rosenfeld, and M. L. Stevenson

Lawrence Radiation Laboratory and Department of Physics, University of California, Berkeley, California

(Received August 14, 1961)

The existence of a heavy neutral meson with $T=0$ and $J=1^-$ was predicted by Nambu¹ in an attempt to explain the electromagnetic form factors

$$|Q|=0: \pi^+\pi^-\pi^0 \quad (800 \times 4 \text{ combinations}), \quad (4)$$

$$|Q|=1: \pi^\pm\pi^\pm\pi^\pm \quad (800 \times 4 \text{ combinations}), \quad (4')$$

PHASE SPACE
 ω MESON
 $\omega \rightarrow \pi^+\pi^-\pi^0$
 $M(\pi^+\pi^-\pi^0)$




FIG. 1. Number of pion triplets versus effective mass (M_3) of the triplets for reaction $\bar{p}+p \rightarrow 2\pi^+ + 2\pi^- + \pi^0$. (A) is the distribution for the combination (4'), $|Q|=1$; (B) is for the combination (4''), $|Q|=2$; and (C) for (4), $Q=0$, with 3200, 1600, and 3200 triplets, respectively. Full width of one interval is 20 Mev. In (D), the combined distributions (A) and (B) (shaded area) are contrasted with distribution (C) (heavy line).

(2). The missing-mass distributions in the two





FIG. 2. (A) M_3 spectrum of the pion triplets in the combined distributions 1(A) and 1(B), with the smooth curve subtracted. (B) M_3 spectrum of the neutral pion triplets in distribution 1(C), again with the smooth background subtracted; a resonance curve is drawn through the peak at 787 Mev with $\Gamma/2=15$ Mev. The error flags are \sqrt{N} , where N is the total number of triplets per 20-Mev interval before subtraction of the smooth background curve.

PHASE SPACE
 SUBTRACTED
 ω
 INVARIANT MASS
 $\pi^+\pi^-\pi^0$

SIMPLE ELECTROSTATIC ACCELERATOR

USED BY COCKROFT & WALTON — ARTIFICIAL RADIOACTIVITY



ELECTRIC FIELD

$$\vec{F} = q \vec{E}$$

CHARGE ON PARTICLE

$$|\vec{E}| = \frac{V}{d}$$

ENERGY GAINED BY CHARGED PARTICLE



$$E_{\text{Acc}} = Fd = qV$$

• TWO SHORTCOMINGS:


— GENERATING HIGH VOLTAGE

— INSULATING BEYOND $\sim 100 \text{ kV}$ (100 keV)

Cockcroft-Walton Generator



VAN DE GRAAFF



• TRANSPORT CHARGE

Q

TO TERMINAL OF CAPACITANCE

C

$$V = \frac{Q}{C}$$

• LIMITATION $\sim 12 \text{ MV}$

\rightarrow VOLTAGE BREAKDOWN


\rightarrow NOT ENOUGH TO

RESOLVE PROTONS

IN THE NUCLEUS

$\sim 12 \text{ MeV}$

TANDEM VAN DE GRAFF



- USE VOLTAGE ON TERMINAL TWICE
- ACCELERATE -VE IONS UP TO TERMINAL
- STRIP OFF TWO ELECTRONS INSIDE TERMINAL



— ACCELERATE AWAY

- 40 MeV CHALK RIVER HAD LARGE TANDEM


VAN DE GRAAFF
IN PRESSURE TANK

ACCELERATED
BEAM

BENDING
MAGNET

1960'S
PHYSICIST

BEAMS TO
EXPERIMENTS




LINEAR ACCELERATOR (LINAC)



TORONTO USED TO HAVE 40 MeV LINAC

• INVENTED BY WIDEROE

RADIO FREQUENCY ω
Generator



• USE SAME RELATIVELY SMALL VOLTAGE IN MANY STEPS — REACH EQUIVALENT HIGH VOLTAGE



- FIELD ZERO INSIDE DRIFT TUBES
- PARTICLE MOVES ONE GAP \rightarrow NEXT, IN TIME E-FIELD REVERSES
- PARTICLES ACCELERATING \rightarrow LENGTH OF DRIFT TUBES INCREASES
- \rightarrow NON RELATIVISTIC

- PARTICLE ENTERING DRIFT TUBE n , ENERGY $n \cdot eV$
↗ VOLTAGE
- NON-RELATIVISTIC
GAPS TRAVERSED ACROSS
KINETIC ENERGY GAP

$$T = \frac{1}{2} m v^2$$

$$v = \left(\frac{2 \cdot n eV}{m} \right)^{\frac{1}{2}} \quad \left(\frac{2T}{m} \right)^{\frac{1}{2}}$$

- THIS VELOCITY TAKES PARTICLE THRU DRIFT TUBE OF LENGTH L_n IN TIME FIELD TAKES TO REVERSE

$$t_n = L_n / v$$

- FREQUENCY OF RADIO FREQUENCY OSCILLATOR

f (Hz) HAS REVERSAL TIME $\frac{1}{2f}$

$$L_n = \frac{1}{2f} \left(\frac{2 n eV}{m} \right)^{\frac{1}{2}} \rightarrow L_n \propto \sqrt{m}$$

NUMERICAL VALUES

$$L_n = \frac{1}{2f} \cdot v_n$$

TYPICALLY $v_n = 0.5c$; $f = 7 \text{ MHz}$ $\rightarrow L_n = 10.7 \text{ m}$

- LOW RADIO FREQUENCY LEADS TO VERY LONG STRUCTURES

- PRACTICALLY NEED HIGH RADIO FREQUENCIES

KLYSTRONS $\rightarrow 100 \text{ MHz} \rightarrow 10 \text{ GHz}$


- THIS WIDEROE STRUCTURE IS OBSOLETE

 - \rightarrow VERY INEFFICIENT




 - \rightarrow RADIATION LOSS

PHASE STABILITY IN LINAC

- TO MAINTAIN PRECISE SYNCHRONISM BETWEEN PARTICLE MOTION & RF OSCILLATOR SEEMS DIFFICULT → NOT SO




- PARTICLE **A** CROSSES GAP1 PHASE ϕ IN STEP WITH VOLTAGE
- GAP2 - SAME VOLTAGE PHASE - AGAIN ACCELERATED
- PARTICLE **B** ARRIVE **LATE**, VOLTAGE **HIGHER**
ACCELERATED **MORE** ARRIVES AT GAP2 **EARLIER**




- PARTICLE C ARRIVES EARLIER AT GAP1
 - VOLTAGE LOWER, ACCELERATED LESS
 - ARRIVES LATER IN PHASE AT GAP2
- B AND C CONVERGE IN PHASE WITH A
- NO NEED TO START WITH PARTICLES ALL IN PHASE WITH RADIO FREQUENCY OSCILLATOR

RADIO FREQUENCY POWER GENERATION




2 CAVITY KLYSTRON OSCILLATOR

ALVAREZ LINAC STRUCTURE



CONDUCTING ENCLOSURE

DRIIFT TUBES



rf generator

DRIIFT TUBES

RADIO FREQUENCY INPUT

- WIDERDE STRUCTURE VERY INEFFICIENT — RADIO FREQUENCY RADIATION LOSS

- ALVAREZ STRUCTURE — RESONANT CAVITY LIKE KLYSTRON

- USED FOR PROTON SYNCHROTRON INJECTOR
100 MeV → 100 MHz


- HIGH ENERGY ELECTRON ACCELERATORS

40 GeV - 500 GeV GHz
RF



SLAC – 50 GeV Electron LINAC

CIRCULAR ACCELERATOR




- REUSE ACCELERATING VOLTAGE UNTIL REACH VERY HIGH ENERGY

PARTICLE BENDING IN MAGNETIC FIELD

$$\vec{F} = q \left(\vec{E} + \frac{1}{c} \vec{v} \times \vec{B} \right) \quad \text{LORENTZ}$$

- FORCE FROM MAGNETIC FIELD NORMAL TO PARTICLE TRAJECTORY



- FOR NO ELECTRIC FIELD & B FIELD NORMAL TO PAGE

$$F = q \frac{v}{c} B \sin \theta \quad \leftarrow 90^\circ = 1 \quad \rightarrow F = q \frac{vB}{c}$$

- FOR A PARTICLE MOVING IN A CIRCLE OF RADIUS ρ

$$\text{CENTRIPETAL FORCE} = \text{LORENTZ FORCE}$$

CIRCULAR ACCELERATORS

- AT PRESENT PARTICLE PHYSICS DOMINATED BY

CIRCULAR ACCELERATORS

ELECTRONS

CESR

PEP II

KEK

LEP

PROTONS

SPS @ CERN

TEVATRON @ FERMILAB

AGS

LARGE HADRON COLLIDER

- MOST EFFICIENT & COMPACT WAY OF GETTING TO HIGH ENERGY - UNTIL SYNCHROTRON RADIATION BECOMES IMPORTANT

CENTRIPETAL FORCE = LORENTZ FORCE

$$\frac{\gamma m v^2}{\rho} = \frac{v B q}{c} \Rightarrow \rho = \frac{p \cdot c}{B q}$$

BENDING RADIUS IN GAUSSIAN UNITS

• ACCELERATOR BUILDERS USE m, VOLT, TESLA

$$pc = p \cdot B \cdot q$$

\nearrow eSV/c \nearrow m \nearrow Gauss

VOLT = 300 STATVOLT / 300
 TESLA = 10^4 GAUSS
 m = 10^2 cm

$$pc \left[\frac{V}{c} \times 300 \right] = p \left[m \times 10^2 \right] B \left[T \times 10^4 \right] e$$

$$pc [GeV/c] = 0.3 p [m] B [T]$$

$$p [m] = \frac{p [GeV]}{0.3 B [T]}$$

$$\phi[\text{sv}]c = \rho(\text{cm}) B(\text{g})$$

$$\phi\left[\frac{\text{V}}{300}\right] \cdot c = \rho(m \times 10^2) B(T \times 10^4)$$

$$\rho c = \rho[m] B[\tau] \times 10^6 \times 3 \times 10^2$$

$$\phi[\text{eV}] = \rho[m] B[\tau] \times 3 \times 10^8$$

$$\phi[\text{GeV} \times 10^9] = \rho[m] B[\tau] \times 3 \times 10^8$$


$$\rho c[\text{GeV}] = \rho \underset{200\text{m}}{[m]} \underset{87}{B[\tau]} \times 0.3$$

$$\rho c = 480 \text{ GeV} = 480 \times 10^9 \text{ eV}$$

$$\rho c[\text{eV}] = \rho(\text{cm}) \times B(\text{g})$$

$$= 200 \times 100 \times 8 \times 10000 = 1.6 \times 10^9 \text{ eSV}$$

$$= 480 \times 10^9 \text{ eV}$$



CENTRIPETAL FORCE = LORENTZ FORCE
FOR AN ORBIT OF RADIUS r

$$\frac{mv^2}{r} = q \frac{v \cdot B}{c}$$

$$\frac{v}{r} = \frac{qB}{mc} = \text{CONSTANT}$$

$$\text{TIME FOR ORBIT} = 2\pi r / v$$

$$\text{ORBITAL FREQUENCY} = v / 2\pi r$$

IF RADIO FREQUENCY f = ORBITAL FREQUENCY

CONTINUOUS ACCELERATION

CONTINUOUS ACCELERATION
RADIO FREQUENCY = ORBITAL FREQUENCY

$$f = \frac{v}{2\pi r} = \frac{1}{2\pi} \frac{qv}{m} \frac{B}{c} = \text{CONSTANT}$$

CYCLOTRON FREQUENCY

↳ DOES NOT DEPEND ON RADIUS
OF ORBIT

- PARTICLE STARTS AT SOURCE CLOSE TO CENTRE OF MACHINE
- SPIRALS OUT CONTINUOUSLY GAINING ENERGY FROM RESONANT RF.

THINK AGAIN ABOUT WHY A CYCLOTRON WORKS

$$F_c = F_L$$

$$\frac{mv}{r} = \frac{q \cdot B}{mc} = k$$

$$\frac{v}{r} = \text{CONSTANT} = \text{FREQUENCY}$$

AS r INCREASES, v INCREASES $\rightarrow \frac{v}{r} = \text{CONSTANT}$
FOR A RELATIVISTIC PARTICLE $v = c = \text{CONSTANT}$

$$\therefore \frac{v}{r} = \frac{c}{r} \neq \text{CONSTANT}$$

ELECTRON CYCLOTRON

"MICROTRON"

ELECTRON IS RELATIVISTIC
FOR $E \sim 500 \text{ keV}$



↓ ORBITS INCREASE
IN RADIUS
DURING
ACCELERATION

ANOTHER RELATIVISTIC EFFECT

$$f = \frac{1}{2\pi} \frac{q}{m} \frac{B}{c} = \text{RF FREQUENCY} = \text{ORBITAL FREQUENCY}$$


- AS PARTICLES ACCELERATE, TOTAL RELATIVISTIC ENERGY BECOMES \approx MASS ENERGY
- IN THIS SITUATION $m \rightarrow m \gamma$ Lorentz Boost

$$f = \frac{1}{2\pi} \frac{q}{\gamma m} \frac{B}{c}$$

DURING ACCELERATION γ INCREASES & RESONANCE CONDITION FAILS

- INCREASE B SYNCHROTRON
- DECREASE RF FREQUENCY SYNCHROCYCLOTRON

SYNCHROTRON - CONSTANT RADIUS ORBIT



DUE TO MAGNETS ONLY AROUND ORBIT
CAN BE MADE VERY LARGE \rightarrow HIGHEST
ENERGIES

SYNCHROTRON

AS USUAL

$$f = \frac{1}{2\pi} \frac{q}{m} \frac{1}{\gamma} \frac{B}{C}$$

EQUAL FOR
RESONANCE

IN RELATIVISTIC SITUATION ORBITAL PERIOD $\frac{2\pi R}{C}$
SO ORBITAL FREQUENCY $C/2\pi R$

CONDITION FOR CONSTANT ACCELERATION


RF FREQUENCY = INTEGER \times ORBITAL FREQUENCY

$$\frac{1}{2\pi} \frac{q}{m} \frac{1}{\gamma} \frac{B}{C} = \frac{C}{2\pi R} \cdot n \quad \leftarrow \text{HARMONIC NUMBER}$$

SINCE $v \approx C \quad \propto \quad p = m\gamma C \quad \rightarrow \quad p \sim m\gamma C$

$$\frac{qB}{p} = \frac{nC}{R}$$
$$R = \frac{mcp}{qB}$$

AS ACCELERATION
PROCEEDS p INCREASES
 $\therefore B$ INCREASES FOR
CONSTANT R




FOR A FIXED B_{MAX}

$$R = \frac{c p}{q B} \Rightarrow R_{MACHINE} \propto p_{MAX}$$

HIGHER ENERGIES \Rightarrow LARGER MACHINES

ELECTRON VERSUS PROTON SYNCHROTRON

- ELECTRONS ACCELERATED AROUND CIRCULAR ORBIT \rightarrow RADIATE



$$\text{ENERGY LOSS} \propto \frac{4\pi e^2}{R} \left(\frac{E}{mc^2} \right)^4$$

$$\frac{\Delta E (\text{PROTON})}{\Delta E (\text{ELECTRON})} = \left(\frac{m_e}{m_p} \right)^4 \approx 10^{-13}$$


THIS IS WHY ELECTRONS IN CERN TUNNEL GO TO 50 GeV WHILE PROTONS TO 7000 GeV

LIMITED BY BENDING
MAGNET




This machine is just a model for a bigger one, of course

1931
 10^4 Volts




Scanned at the American Institute of Physics




This machine is just a model for a bigger one, of course

1953
 10^9 Volts

Cosmotron Coil Built by HYRE ELECTRIC CO. CHICAGO




1932
 10^6 Volts



This machine is just a model for a bigger one, of course

1960
 10^{10} Volts




Livingston and Lawrence with the magnet of the “27-inch” (later “37-inch”) cyclotron on which most of Berkeley’s 1930s nuclear physics was performed.
Lab wear was different then!

THE 184-INCH SYNCHROCYCLOTRON



The Berkeley 184" was begun in 1939 as a classical cyclotron, to be operated with $V_{rf} = 1$ MV, but WWII interrupted rf installation and it was used to test mass spectrographic separation of uranium isotopes. **FM rf was installed in 1946**, yielding **190 MeV d+** (700 MeV p in 1959).

PET Medical Cyclotron





TRIUMF (Vancouver) 500 MeV Cyclotron






ACCEL

250 MeV Superconducting Proton Cyclotron



Superconducting Coil



The present study is partially supported by the REIMEI Research Resources of Japan Atomic Energy Research Institute.

You can have your own cyclotron – from Mitsubishi



Alors, c'est fini!
Et maintenant?

DC HIGH-VOLTAGE ACCELERATORS – TANDEM VAN DE GRAAFFS




Yale 22-MV tandem.



Daresbury folded tandem
(20 MV in a 230-ft tower).



The ISAC 150-keV/u RFQ linac



500 keV electron LINAC for Cancer Therapy

