


# **Hard Scattering in Hadron-Hadron Collisions: Physics and Anatomy**

## **Section 4: Production & Identification of Jets**

- 1. Definitions of Basic Physics Processes**
- 2. Anatomy of a Jet**
- 3. Jet-Finding Algorithms**
- 4. Resolutions and Efficiencies**
- 5. Heavy Quark Tagging**
- 6. Example: Quark Substructure**

# Definitional Issues

- **Confinement in QCD ensures that high  $P_T$  quarks & gluons undergo**
  - **Fragmentation -- ie, dissociation into a “jet” of coloured partons**
  - **Hadronization -- ie, the partons form colourless, observable hadrons**
- **Study of jets motivated by**
  - **Understanding QCD**
  - **Studying of heavy quarks**
    - > b/c quarks that fragment & hadronize before decay
    - > Top quarks that decay before fragmentation/hadronization
  - **Searching for new interactions that couple to quarks/gluons**
  - **Backgrounds to  $e, \mu, \gamma$  &  $\tau$**





# Fundamentals of Jet Physics

- Basic production mechanism in pQCD starts with

$$\sigma = \sum_{\substack{\text{partons } i \\ \text{colour } j}} C_{ij} \int_0^1 d\tau \int_{\tau}^1 \frac{dx_1}{\tau} [f_1(x_1) f_2(\tau/x_1)] \hat{\sigma}(\tau s)$$

- Leading-order (LO) diagrams already complex




Process	$\bar{\Sigma}  \mathcal{M} ^2 / g^4$	Numerical value for 90°
$qq' \rightarrow qq'$	$\frac{4}{9} \frac{s^2 + u^2}{t^2}$	2.22
$q\bar{q}' \rightarrow q\bar{q}'$	$\frac{4}{9} \frac{s^2 + u^2}{t^2}$	2.22
$qq \rightarrow qq$	$\frac{4}{9} \left( \frac{s^2 + u^2}{t^2} + \frac{s^2 + t^2}{u^2} \right) - \frac{8}{27} \frac{s^2}{ut}$	3.26
$q\bar{q} \rightarrow q'\bar{q}'$	$\frac{4}{9} \frac{t^2 + u^2}{s^2}$	2.22
$q\bar{q} \rightarrow q\bar{q}$	$\frac{4}{9} \left( \frac{s^2 + u^2}{t^2} + \frac{t^2 + u^2}{s^2} \right) - \frac{8}{27} \frac{u^2}{st}$	2.59
$q\bar{q} \rightarrow gg$	$\frac{32}{27} \frac{t^2 + u^2}{tu} - \frac{8}{3} \frac{t^2 + u^2}{s^2}$	1.04
$gg \rightarrow q\bar{q}$	$\frac{1}{6} \frac{t^2 + u^2}{tu} - \frac{3}{8} \frac{t^2 + u^2}{s^2}$	0.15
$gq \rightarrow gq$	$\frac{4}{9} \frac{s^2 + u^2}{su} + \frac{u^2 + s^2}{t^2}$	6.11
$gg \rightarrow gg$	$\frac{9}{2} \left( 3 - \frac{tu}{s^2} - \frac{su}{t^2} - \frac{st}{u^2} \right)$	30.4

# What Have We Learned?


- **Definition of jets critical**
  - Much evolution in algorithms
  - Driven in large measure by theoretical considerations
- **Calibration of jets requires data-driven techniques**
  - Developed several techniques to calibrate *in situ*
  - Still “work in progress”
- **Approach to jet-finding and calibration driven by physics**
  - Best example is comparison between
    - > QCD tests
    - > Reconstruction of heavy objects (top and Higgs)

$$\Psi(r) \equiv \frac{1}{N_{jet}} \sum_{jets} \frac{P_T(0,r)}{P_T(0,R)}$$



D. Acosta et al. (CDF), Phys. Rev. D 71, 112002 (2005)


- **Need data to understand jets as backgrounds**
  - Examples include
    - > Lepton ID
    - > MET measurement
    - > Heavy quark tagging
  - Use to “calibrate” MC/simulation
- **Bottom line: SM Picture of QCD works well**




# LHC Lessons

## ■ LHC studies have reproduced many of these observations

- However, much higher jet momenta
  - > Jets with  $p_T > 2$  TeV
- Focus has been on searches
  - > Looking for resonances in dijet mass
  - > Sensitive to excited quarks




ATLAS Collaboration, Phys. Rev. D 83, 052003 (2011)



ATLAS-CONF-2011-096, 18 July 2011

# Highest Dijet Mass Event



# Jet Anatomy

## ■ A jet arises from 2 different physical phenomena

- Happen at different energy scales
  - > Fragmentation of initial parton
    - QCD radiation of a coloured object
    - Creates a “cluster” of coloured partons
    - In principle, not independent of rest of event
    - Energy scale  $\gg 1$  GeV
  - > Hadronization of “cluster”
    - Formation of colourless objects -- mesons & baryons
    - Responsible for the real observables
    - Energy scale  $\sim 1$  GeV

## ■ Have to worry about

- What defines a jet (algorithm)?
- What its properties are (recombination scheme)?

## ■ First, tackle easiest part: What is a jet’s observable properties?

- Assume you have a collection of final state mass-less “particles” detected in calorimeter towers  $i$

$$\vec{p}_J \equiv (E^J, p_x^J, p_y^J, p_z^J) \equiv \sum_i (E^i, p_x^i, p_y^i, p_z^i)$$

$$p_T^J \equiv \sqrt{(p_x^J)^2 + (p_y^J)^2}$$


$$M^J \equiv \sqrt{(E^J)^2 - (p^J)^2}$$

$$y^J \equiv \frac{1}{2} \ln \frac{E^J + p_z^J}{E^J - p_z^J}$$

$$\varphi^J \equiv \tan^{-1} \frac{p_y^J}{p_x^J}$$

## – Advantages:

- > Clear Lorentz behaviour
- > Avoids use of  $E_T$  which has ill-defined definition
- > Can generalize to “cells”, towers, charged particles, etc.







# A Real Jet Event

Tower  $E_T > 0.5$  GeV


$K_T$  D=0.7: Raw  $P_T^{\text{JET}}$



**CDF RUN II**  
Run 163064  
Event 6753986



# An LHC CMS Jet Event



Highest dijet  
Mass event  
Observed by  
CMS at 7 TeV

CMS Collaboration, hep-ex/1107.4771

# Parton Shower Evolution

- Start with a parton (q/g) with virtuality  $\mu^2$




- Probability of emission with daughter carrying  $z$  fraction of parent momentum

$$d^2 P_a(z, \mu^2) = \frac{d\mu^2}{\mu^2} \frac{\alpha_s}{2\pi} P_{a \rightarrow bc}(z) dz$$

- Order these using Sudakov factor, relating  $\mu^2 \sim Q^2$

$$P_a^{no}(Q_{\max}^2, Q^2) = \exp\left(-\int_{Q^2}^{Q_{\max}^2} dQ' \int_{z_{\min}}^{z_{\max}} dz' P_a(z', Q'^2)\right)$$


- Deal with infrared & collinear divergences
  - > Define minimum  $\mu - \mu_0$
- Ensure colour coherence of multiple emissions
  - > Typically do this by angular ordering, selective vetoing, etc.
  - > Must be respected when hadronization is performed



# Hadronization of Showers



## Hadronization is then performed

- Invoke “parton-hadron duality”
- Several models
  - > String fragmentation (eg., PYTHIA)
  - > Cluster fragmentation (eg. HERWIG)



- Have various parameters that need to be tuned to data
  - > Best constraints from LEP
    - Tevatron results confirm these, but don't really add much power
    - Challenging to measure without significant systematic uncertainties
  - > Remains a source of systematic uncertainty

OPAL, Eur. Phys. J C16, 185 (2000)



# Jet Algorithms





- **Jet clustering algorithms have evolved over the last 30 years**
  - **Goals of any algorithm can be divided into**
    - > **Theoretically motivated:**
      - **Fully specified**
      - **Detector independent**
      - **Theoretically well-behaved**
      - **Order independent**
    - > **Experimentally motivated:**
      - **Fully specified**
      - **Detector independent**
      - **Optimal resolution and efficiency**
      - **Ease of calibration**
      - **Computationally efficient**
- **Various efforts to develop consistent frameworks**
  - **Snowmass Accord (1990)**
  - **Les Houches Accord (1999)**
- **Raz Alon (see talk below) has done a nice job of summarizing current Jet Algorithm codes**
  - **Key observations:**
    - > **In principle, prefer some algorithms over others**
      - **Seedless cone-based algorithms**
      - **$K_T$  algorithms**
    - > **Computational efficiency is a concern in some cases**
      - **But largely an issue of optimization**
    - > **Selection of “best” algorithm requires evaluation of ultimate systematic uncertainties**
      - **Need data, as certain choices will depend on performance of calorimeter**
      - **Example is noise and pileup**
  - **Good news is that we are not limited by lack of ideas**

R. Alon, <http://indico.cern.ch/conferenceDisplay.py?confId=52628>

# Clustering Effects




## ■ Illustrate by one example (from ATLAS studies)

- Compare results of several different algorithms
  - >  $K_T$  with  $R=1$
  - > Angular-ordering (Cam/Aachen)
  - > SISConc
  - > Anti- $K_T$
- Things to be concerned about
  - > Cluster sizes determined by data will present challenges to calibrate
  - > Cluster merging/splitting will continue to be a challenge
  - > Optimization of resolution/systematic uncertainties will require effort
- Things not to worry about
  - > Angular resolution (though need to check for any biases)!



# Jet Finding Efficiencies

- Efficiency of **finding** jets limited primarily by two effects:
  - Detector energy response & resolution
  - Physical size of jets
    - > For cone algorithms, these two compete with each other
- Further complicated by the fact that jets are produced with sharply falling spectrum
  - Means that efficiencies become an issue already at the trigger level
  - Manage these at Tevatron & LHC with variety of triggers
    - > Prescale lower-energy jet triggers ( $\sim 10^3$ )
    - > Lower energy jets used primarily for
      - Background studies
      - Calibration





# Jet Energy Resolutions

- MC & simulation give estimates of energy resolution

- Resolution is determined primarily by convolution of
  - > Intrinsic calorimeter response
  - > Jet fragmentation & hadronization effects
  - > Jet algorithm + pileup + ....
- In reality, need to measure the resolution in data

- Four *in situ* measurements of resolution developed at Tevatron


- $\gamma$ +jet balancing
- W to qq in top quark decays
- Dijet balancing (more of a constraint than anything else)
- Z to bb decays
  - > Require two jets, each with secondary vertex b-tag
    - Possible due to L2 vertex trigger

- CDF: Taking the FWHM  $\sim 25 \text{ GeV}/c^2$ , obtain

$$\sigma_Z \sim 12\% M_Z$$

$$\Rightarrow \frac{\sigma}{P_T^J} \sim 17\%$$

- Or about 50% more than intrinsic energy resolution of calorimeter






# Jet Energy Calibration

## ■ To calibrate jet energy scale:



### – 1. Determine intrinsic response to particles

- > Combination of in situ measurements & test beam data



### – 2. Dijet balancing to get uniform $\eta$ response

- > Primarily dijet data
- > “Tune” MC and simulation





### – 3. Determine absolute response to “particle jet”

- > Define particle jet as all real particles in cone of jet
- > Account for calorimeter nonlinearity, cracks, etc.

### – 4. Take into account “out-of-cone” effects, multiple interactions

- > Use combination of MC and data




# Final Steps in Energy Calibration


## ■ Cross check using, for example,

- Z+jet &  $\gamma$ +jet balancing
- Dijet balancing
- W  $\rightarrow$  jj in ttbar events


W  $\rightarrow$  jj



Z-jet Balancing




$\gamma$ -jet Balancing



## ■ Estimate systematic uncertainties



- Estimate each source independently
- Struggle with the fact that we cannot measure high  $P_T$  jet response



# ATLAS Jet Calibration

## ■ Used similar techniques


- Z+jet &  $\gamma$ +jet balancing
- Dijet balancing
- Also extended and tested other techniques
- Have 2 calibration schemes
  - > EM+JES
  - > LCW+JES



ATLAS Collaboration, Eur. Phys. J. C (2015) 75:17, [arXiv:1406.0076](https://arxiv.org/abs/1406.0076)

# Production Cross Sections

- CDF analysis of  $\sim 1.13 \text{ fb}^{-1}$  of jet data
  - Used mid-point algorithm with  $R=0.7$ ,  $f_{\text{merge}}=0.75$
  - Data is scaled in plot to avoid overlapping
- Provide a strong test of QCD
  - Theoretically “clean” to model
  - Compare with NLO calculations
    - > Fill in details!
  - Generally a trend of small excess of events at higher  $P_T$
  - Not statistically significant given systematic uncertainties




# Total Jet Production Rates

## ■ An “Exercise to Reader” – what is total cross section?

- To answer this question
  - > Fit the spectrum in each  $y$  bin to power law using ROOT
  - > Use fit to extrapolate over various  $P_T$  ranges
    - Was lazy, only did the first four bins
  - > Generally, differential cross section falls with  $(P_T)^{-6}$ 
    - And gets a little steeper as  $P_T$  increases
    - Means that higher  $P_T$  jets tend to be more central

## ■ Note large cross section at low $P_T$

- This is the source of backgrounds to other objects
- Also note that these are quite uncertain given the extrapolation!
  - > Eg., just changing range of fit
    - $\Delta\sigma(P_T > 10) \sim 30\%$





Cross Section (in nb)

	PT > 62 GeV	PT > 30 GeV	PT > 10 GeV
$ y  < 0.1$	122	5,600	1,800,000
$0.1 <  y  < 0.7$	111	5,600	2,000,000
$0.7 <  y  < 1.1$	96	6,100	3,000,000
$1.1 <  y  < 1.6$	93	8,900	8,900,000
	422	26,200	15,700,000

Note: Another ~5-10% in rapidity interval  $1.6 < |y| < 2.1$

# Heavy Quark Jets


- **Heavy quarks (b/c) also manifest themselves as jets**
  - **Different fragmentation process**
  - **Different hadronization**
    - > Result in kinematics that differ from light quark & gluon jets
  - **“rich” in  $\nu$ 's and charged leptons**
    - > Used for identification
    - > But also affect efficiency and energy resolution
  - **Relatively long lifetimes allow for tagging using secondary vertices**
    - > Become “standard” technique
- **Bottom quarks have been particularly important**
  - **Essential for top quark studies**
  - **Result in unique capabilities at hadron colliders**
    - > Good example is  $B_s$  studies



# Heavy Flavour Tagging

- Heavy flavour tagging has been essential tool at Tevatron
  - Top quark search
  - Search for Higgs
  - Studies of bottom/charm production
- Two methods developed
  - Semileptonic tagging
    - > 20% of  $b'$ 's decay inclusively to  $\mu$  or  $e$ 
      - Another 20% have leptons from charm decay
    - > Challenge is purity of tagging scheme
      - CDF couldn't get fake rates below about 3-4%
  - Secondary vertex tagging most powerful


- Basic strategy is to use well-measured tracks
  - Select those with large impact parameter
    - > Typically reconstruct average primary beam position in  $(x,y)$
  - Require 2+ tracks with impact parameter  $> 2s$  and high quality
    - > Attempt to create a secondary vertex
    - > If successful, see if secondary vertex is sufficiently far from primary
      - Tag when secondary vtx found
      - Also “fake tag” when tag found, but in wrong direction






# Tagging Efficiencies


- **Tagging efficiency difficult to model via simulation**
  - Requires excellent knowledge of tracking resolution & efficiency
  - **Strategy:**
    - > Measure efficiency and “mistag” rates in data
      - Inclusive electrons and muons
        - Estimate b quark fraction
      - Tag fully reconstructed Bs
    - > Compare with simulation & compute a scale factor
      - $SF = \epsilon_{\text{data}} / \epsilon_{\text{MC}} \sim 0.95 \pm 0.05$  for “tight” SECVTX



**SecVtx Tag Efficiency for Top b-Jets**



**SecVtx Tag Efficiency for Top b-Jets**





# Tagging Fake Rates

- **B tagging fake rates measured from data**
  - Take samples of dijet data, and then create a “fake matrix”
    - > Function of 6 variables
    - > Measure both +ve and -ve tag rates for “taggable jets”
      - Use -ve tag rates as mistag rate
    - > Apply mistag rate to the jets in data sample before tagging

Variable	Bin Edges
Jet $E_T$ (GeV)	0, 15, 22, 30, 40, 60, 90, 130, 1000
Num Tracks/jet	0, 1, 2, 3, 4, 5, 6, 7, 8, 10, 13, 100
$ \eta_{\text{jet}} $	0.0, 0.4, 0.8, 1.1, 2.4
$n_{\text{PrimaryVertex}}$	1 – 6
$\sum E_t$ (GeV)	0, 80, 140, 220, 1000
$z_{\text{prim}}$ (cm)	-25, -10, 10, 25

Table 1: Variables and binning used in the mistag matrices.



# Example: Quark Substructure

## ■ Search for quark substructure a long-standing tradition at high energies

- Eichten, Lane & Peskin
  - > PRL 50, 811 (1983)
- Introduced “contact term”  $\Lambda_C$
- CDF obliged in 1996
  - >  $\Lambda_C \sim 1.6$  TeV





FIG. 3. The jet production cross section (in picobarns/gigaelectronvolt) at rapidity  $y = 0$  vs transverse momentum at  $\sqrt{s} = 2$  TeV in (a)  $\bar{p}p$  collisions and (b)  $pp$  collisions for various  $\Lambda$  (in teraelectronvolts). The solid and dashed lines in (b) refer, respectively, to the plus and minus signs in Eq. (5). As a result of a cancellation near  $y = 0$ , the interference is negligible in (a).




## ■ Later shown to be described by different PDF behaviour at large x

# More Sensitive Study

- Employ angular distribution in dijet scattering

$$\chi \equiv \exp|\eta_1 - \eta_2|$$


- Look at this as a function of dijet invariant mass
  - > 100 GeV mass bins
- More sensitive to  $\Lambda_C$ 
  - > Less sensitive to PDFs
  - >  $\Lambda_C > 2.4$  TeV at 95% CL



# And Even More Sensitive!

- **ATLAS has further improved sensitivity**
  - Look at fraction of centrally produced jets relative to larger angular range
  - See behaviour as dijet mass increases
  - Expect QCD background to have flat ratio
- **More sensitive to  $\Lambda_C$** 
  - $\Lambda_C > 9.5$  TeV at 95% CL

ATLAS Collaboration, New J. Phys. 13,053004 (2011)



$$F_{\chi}([m_{jj}^{\max} + m_{jj}^{\max}] / 2) \equiv \frac{N_{\text{events}}(|y^*| < 0.6, m_{jj}^{\min}, m_{jj}^{\max})}{N_{\text{events}}(|y^*| < 1.7, m_{jj}^{\min}, m_{jj}^{\max})}$$

$$y^* \equiv \frac{1}{2}(y_1 - y_2)$$