CDF/ANAL/CDF/PUBLIC/5920

A Summary of the Treatment of Systematics in the CDF Run I Top Cross Section Measurement

Pekka K. Sinervo

University of Toronto

13 April 2002

1 Introduction

The CDF collaboration has published a final measurement of the top quark production cross section using data collected during 1992-1995 (Run I) [1]. This measurement is basically a counting experiment, where one relates the production cross section, $\sigma_{t\bar{t}}$, to the number of candidate events, $N_{t\bar{t}}$, using the formula

$$\sigma_{t\bar{t}} = \frac{N_{t\bar{t}} - N_{bkg}}{\epsilon\mathcal{L}}, \tag{1}$$

where N_{bkg} is the number of background events in the candidate sample, ϵ is the efficiency for identifying a $t\bar{t}$ event as a member of the candidate sample, and $\mathcal{L}$ is the integrated luminosity for the experiment.

The "random" uncertainties associated with this measurement come from the Poisson fluctuations in $N_{t\bar{t}}$. This note summarizes the treatment of uncertainties in this measurement in a context where a number of channels were combined together to give a final Run I result.

The primary reference to the measurement discussed here [3] provides a very nice capsule overview of how the final result was determined. Details on the cross section measurements in the individual channels are provided in a series of notes referenced therein.

2 Definition of Uncertainties

We conventionally divide sources of uncertainty into two classes, "random" or "statistical" uncertainties, and "systematic" uncertainties. Random uncertainies are those that would normally scale by the total size of the data sample, typically with a $1/\sqrt{N}$ dependence. Systematic uncertainties can be defined in various ways [2], but for this measurement are those that effect a parameter or procedure used in extracting the result from the data.

Statistical uncertainties arise from the random fluctuations in the data sample. In the case of an observation where one counts the number of events, the Poisson fluctuations in

the statistic of relevance, namely the number of observed events, would be the source of the statistical uncertainty. In more complex examples, it is typical to define a likelihood function that describes the data and use the shape of the likelihood function to determine a statistical uncertainty.

Systematic uncertainties, on the other hand, are the result of uncertainties in various parameters or inputs into the measurement that are required to interpret the results, or more generally uncertainties in the experimental conditions or the theoretical model used to interpret the data. These uncertainties can be characterized in a number of ways. There are in principle some sources of uncertainty where the effects scale with the total number of events, but are considered by convention to be systematic uncertainties. There are on the other hand, those sources of uncertainty that do not have any dependence on the size of the data sample. There are at two least reasons why we would wish to make the distinction between these two types of systematic uncertainty:

1. The uncertainties that depend on the event sample size will scale with the total integrated luminosity of the sample, and will therefore be reduced in future running. The other class of uncertainties will not be reduced unless steps are taken to improve the measurement.

2. The two classes of uncertainty are likely to create different correlations between our measurement and measurements made by a different experiment or different technique. This difference should be taken into account when combining two or more measurements.

3 The Total Cross Section Measurement

The CDF collaboration detected pair production of top quarks by selecting events in which there was evidence of the semileptonic or hadronic decay of two top quarks. Events in which both top quarks decays semileptonically, *ie.* $t \to bl\nu_l$, where l denotes either a muon or electron, are by far the cleanest since the backgrounds associated with a dilepton signal are relatively low. It is, however, a small sample due to the approximately 10% semileptonic branching fraction for a single top quark. The largest sample of events arises in the case where both top quarks decay hadronically, *ie.* $t \to bq\bar{q}'$, but this sample also is severely contaminated with background arising from QCD multi-jet production. The sample arising from the case where one top quark decays semileptonically and the other hadronically produces an event signature consisting of a single lepton candidate, missing transverse energy from the neutrino and in principle 4 quark jets. This "lepton+jets" channel is intermediate in size between the dilepton and all hadronic channel, and has moderate background contamination (most of the background arises from $W + b\bar{b} + X$ production).

Thus, the strategy used in the Run I cross section measurement was to select samples of top quark candidates for these three channels, estimate the backgrounds in each sample, and then correct the estimated number of signal events for acceptance and efficiency effects to derive a cross section measurement in each channel. The final step was to combine the measurements from the different channels into a single measurement.

The primary data for this measurement are summarized in Table 1. The method used to combine these data was to perform a maximum likelihood fit, where the likelihood was

Channel	Total Events	Background Rate	Aceptance×Efficiency
Dileptons	9	2.4 ± 0.5	0.0074 ± 0.0008
Lepton+Jets (SVX tags)	29	8.0 ± 1.0	0.039 ± 0.006
Lepton+Jets (SLT tags)	25	13.2 ± 1.2	0.012 ± 0.002
All Hadronic (kinematic)	42.8 ± 17.9		0.055 ± 0.012
All Hadronic (double tag)	36.7 ± 13.7		0.045 ± 0.014

Table 1: The data for the Run I top quark cross section measurement.

parametrized as a function of $\sigma_{t\bar{t}}$.

4 Random Uncertainties

The random uncertainties in this measurement arise from the total number of candidate events, namely 29 observed SVX-tagged events, 25 SLT-tagged events, 9 dilepton events and the observed rate of all-hadronic events above background. The two lepton+jets samples have an overlap of 7 events but this is ignored (as the effect has been shown to be less than 10% in the overall uncertainty of the cross section derived from these two measurements). There are two samples of hadronic candidate events with large backgrounds. The background-subtracted numbers of events for these two samples are 42.8 ± 17.9 and 36.7 ± 13.7, which are assumed to be correlated Gaussian statistics. The correlation in the two all-hadronic event samples was significant and was modelled by including a correlation coefficient $\rho = 0.34$ in the joint probability distribution for the number of observed all-hadronic events in the sample

$$G_{had}(N_{had1}, N_{had2}, \rho, \mu_{had1}, \mu_{had2}), \tag{2}$$

where μ_{had1} and μ_{had2} are the expected mean number of observed events (and are functions of the $\sigma_{t\bar{t}}$). Note that this latter term does not include any background contribution as the event rates N_{had1} and N_{had2} are both numbers of signal events *above* background. The form of this likelihood distribution is shown in Fig. 1.

The uncertainties associated with these numbers are incorporated into the final result through the use of a maximum likelihood fit. In effect, the numbers of dilepton and lepton+jet events are treated as Poisson statistics. In order to combine the "all hadronic" events, the likelihood function included the the two-dimensional Gaussian probability distribution G_{had}. The means μ_{had1} and μ_{had2} of this Gaussian distribution would be given by the total top quark cross section times its branching fraction and acceptances into the two all-hadronic channels. The new likelihood function was minimized with respect to the top quark cross section.

The overall form of the likelihood function is

$$\begin{aligned} L \;=\; & \left[\Pi_i^{svx,slt,dil} P(N_i, \mu_i(\sigma_{t\bar{t}}, ...))\right] \times \\ & G(N_{had1}, N_{had2}, \rho, \mu_{had1}, \mu_{had2}) \;\times\; \Pi_{j=1}^{12} G(X_j, \bar{X}_j, \sigma_j) \end{aligned} \tag{3}$$

where the first two terms represent the statistical uncertainties (the Poisson probability distributions for the observed number of events and the Gaussian probability distribution



Figure 1: The form of the likelihood distribution for the two correlated hadronic channel measurements.

describing the uncertainties in the two all-hadronic rates). The last term represents the 12 sources of systematic uncertainty, discussed below, where G is a one-dimensional Gaussian distribution with the form

$$G(x_o, \sigma_o; \mu) \propto \exp\left(\frac{-(x_o - \mu)^2}{2\sigma_o^2}\right), \tag{4}$$

where x_o and σ_o is the fitted deviation of the parameter suffering the systematic uncertainty and its associated uncertainty.

5 Systematic Uncertainties

The philosophy chosen in this analysis was to estimate the individual effects that created further uncertainty in the resulting top quark cross section. These effects ranged from detector acceptance, trigger efficiency and lepton identification efficiency to uncertainties in the modelling of the top quark interaction (which then affect the measurement by modifying the expected top quark acceptance and efficiency). Overall, there are 12 different sources

of systematic uncertainty, ranging in relative size from about 4% to about 70%. Typically, each of these has been estimated from studies based on either data or Monte Carlo samples (often both), with the goal of identifying a range that corresponds to a 68% confidence level interval. The very largest uncertainty (70%) is associated with a relatively small component of a background source and so does not contribute significantly to the overall uncertainty in $\sigma_{t\bar{t}}$.

Although the sources of these additional uncertainties are quite different, they were incorporated into the final result by treating them as Gaussian statistics with normal distributions characterized by the uncertainty assigned to the central value of a given parameter. With this assumption, they could be treated as being another set of uncertainties that could be included in the likelihood function. With this approach, all of the uncertainties associated with the measurement were accommodated in the likelihood function. The cost of this was that the likelihood was now no longer just a function of the parameter of interest ($\sigma_{t\bar{t}}$), but also of twelve "nuisance" parameters.

The log of the likelihood function was minimized using MINUIT and the uncertainty in the cross section was evaluated by determining by mapping the likelihood function – the cross section was fixed at specific values and the log-likelihood minimized with respect to all the other parameters. The uncertainty in the cross section was then defined to be the one-half unit change in the log-likelihood from its minimum value. The result was

$$\sigma_{t\bar{t}} \;=\; 6.55^{+1.68}_{-1.41}\ \text{pb}, \tag{5}$$

where the uncertainty now includes both statistical and systematic effects.

6 Discussion

The choice to include in the likelihood function all of the systematic uncertainties is one that has been employed in many other analyses within CDF [4]. It has the convenient effect of burying in the likelihood both statistical and systematic uncertainties and providing a formula for combining them. Clearly, this procedure rests on the assumption that one can describe one's uncertainty in the systematic effects with a Gaussian probability distribution. We believe this to be the case, but this depends on the results of numerous cross checks that are reassuring but do not guarantee that all effects have been accounted for.

Of perhaps more philosophical interest is the interpretation of this likelihood function for those wishing to use it to place confidence intervals on the top quark cross section. In principle, a Bayesian statistician could assume a prior distribution for $\sigma_{t\bar{t}}$, multiply the likelihood function by the prior and use the resulting function as the posteriori probability for $\sigma_{t\bar{t}}$.

A frequentist statician would have a more difficult time using the likelihood, as he or she would now have to define the appropriate ensemble that should be used in setting a frequentist limit. In principle, one could model the ensemble as a repetition of the measurement, taking into account changes in procedure and assumptions that result in the variations in the parameters that introduce the "systematic uncertainties." To the extent that each parameter does have a single true value and the experiment provides different estimates of its value when repeated (as modelled by the Gaussian distribution function incorporated

into the likelihood function), a standard frequentist confidence limit calculation could be performed. However, this approach could be computationally very prohibitive. Practical approaches to address this calculation are available [5].

This interpretation is not possible in the context where the systematic uncertainty arises from theoretical considerations that we are unable to constrain from the data. In this case, an alternate approach would be to not include such theoretical effects as systematic uncertainties, but instead to characterize the sensitivity of the final result (and its uncertainty) on variations in the theoretical assumptions. This would more clearly identify such assumptions and allow different experiments to more effectively combine their measurements using a common framework. However, a sensitivity analysis would then have to be provided.

References

[1] *Measurement of the $t\bar{t}$ cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*,
T. Affolder *et al.*, Phys. Rev. D **64**, 032002 (2001).

[2] The CDF Statistics web page section on systematic uncertainties gives several operational definitions and practical examples.

[3] CDF Note 5043, The Top Group (June 1999).

[4] *Measurement of the CP Violation Parameter* $\sin(2\beta)$ *in* $B^0\overline{B^0} \to J/\psi K$ *Decays*,
F. Abe. *et al.*, Phys. Rev. Lett. **81**, 5513-5518 (1998).

Measurement of b quark fragmentation fractions in the production of strange and light B mesons in proton anti-proton collisions at $\sqrt{s} = 1.8$ TeV,
F. Abe *et al.*, Phys. Rev. D. **60**, 092005 (1999).

Measurement of b Quark Fragmentation Fractions in $\bar{p}p$ Collisions at $\sqrt{s} = 1.8$ TeV,
T. Affolder *et al.*, Phys. Rev. Lett. **84**, 1663-1668 (2000).

[5] See, for example, the CDF Statistics web page and notes referenced therein.