A Five Year Plan for CDF

Pekka K. Sinervo
University of Toronto / IPP

1. CDF Physics
2. Run 1 Status
3. Run 2 Proposal
4. Financial and Personnel Requirements

TRIUMF
September 30, 1995

Physics at 2 TeV

$p\bar{p}$ collisions at 1.8 TeV open unique window onto:

- $W^{\pm}$, Z° and γ production
- Perturbative QCD tests
- Heavy (b and c) quark production & spectroscopy
- Top Quark Physics
- EWK tests: M_W, WWZ couplings, Γ_W
- New physics: SUSY, compositeness, $H^{\pm}$, technicolor



Many new vistas open up with very high sensitivity (0.1 $\rightarrow\sim$ 1 fb^{-1})

- Detect $\lesssim$ 0.01 pb cross sections
- Explore higher-order EWK processes
- Extend "mass reach" for new phenomena by $\sim\times 2$
- High statistics b and t physics

CDF Run 1 Physics

CDF collaboration has well established track record

- Operates detector at Fermilab Tevatron Collider
- Collaboration consists of $\sim$ 450 physicists and 35
- Involves groups from five countries

CDF is just finishing a three-year run (1A and 1B)

- In 1992-93, collected $\sim$ 20 pb^{-1} (Run IA)
- Since 1994, have collected $\sim$ 100 pb^{-1}(Run 1B)
  - Run 1B continuing through February 1996
  - Ultimate goal is $\sim$ 150 pb^{-1}

Many new results have already been published

- New measurement of M_W and Γ_W
- Discovery of top
- Wealth of bottom physics

$\boxed{W \text{ Mass Measurement}}$

W production cross section is ~ 20 nb at Tevatron

- Can readily observe $W \to e\nu_e$ and $W \to \mu\nu_\mu$
- Select high E_T electron or muon
- Require significant $\not{E}_T$ to tag neutrino

Can extract M_W from transverse mass distribution

$$M_T = \sqrt{2E_T \not{E}_T (1 - \cos\phi_{l\nu})}$$

- Sensitive to performance of entire calorimeter
- Sensitive to structure functions



Fit $65 < M_T < 100$ GeV to predicted lineshape

- 5718 electron events
- 3268 muon events

Fit to Muon Data

Results of fit to muon data

$$M_W = 80.310 \pm 0.205 \pm 0.130 \text{ GeV/c}^2$$



Fit to Electron Data

Results of fit to electron data

$$M_W = 80.490 \pm 0.145 \pm 0.175 \text{ GeV/c}^2$$



Systematic Uncertainties on W Mass

Control of systematics most challenging aspect

1. Energy and momentum scale (120 and 45 MeV/c^2)
   - Determined using $\Upsilon/\psi \rightarrow \mu^+\mu^-$ and $Z \rightarrow \mu^+\mu^-/e^+e^-$
   - calorimeter scale set by tracking system

2. Resolution function of "detector" ($\sim$ 80 MeV/c^2)
   - Constrained by studies of Z events

3. Uncertainies in "recoil" system physics ($\sim$ 60 MeV/c^2)
   - Limited by statistics on Z sample

4. Structure functions (50 MeV/c^2)
   - Use MRS D_-' as default
   - Check variation caused by other choices

Accounting for correlations in these uncertainties

- Overall systematic uncertainty is 130 MeV/c^2

Summary of W Mass

Combined mass from both channels is

$$M_W = 80.41 \pm 0.18 \text{ GeV/c}^2$$



$\boxed{\text{Dilepton Top Quark Search}}$

$t\bar{t}$ decaying into 2 leptons create unique signature

- Two charged leptons from $W^{\pm}$ decay
- Imbalance in total energy due to undetected neutrinos
- Two jets from bottom quarks

CDF performed a search requiring

- Two clean, high P_T lepton candidates
- Missing transverse energy $\not{E}_T > 25$ GeV
- Require two jets with E_T> 10 GeV and $|\eta| < 2.0$
- Reject specific backgrounds (e.g. Z°)

Electron-Muon Data for Run 1A and 1B

$e\mu$ data: $\Delta\varphi$ (MET, lepton or jet) vs Missing E_t



Summary of Dilepton Search

Observe a total of 6 candidate events

- Observe 1 $\mu^+\mu^-$ event and 5 $e\mu$ events
- Expect 1.3 ± 0.3 events from background

Background	Expected Background Rate
$Z \to \tau^+\tau^-$	0.38 ± 0.07
One or both leptons fake	0.23 ± 0.15
$b\bar{b}$	0.03 ± 0.02
WW, WZ and ZZ	0.21 ± 0.07
Drell-Yan	0.44 ± 0.28
Total	1.3 ± 0.3

Probability of background fluctuation is 3.3×10^{-3}

b Quark Tagging in W+Jet Events

Top observed in lepton+jets channel where we look for

$$t \to W^{+}b \to l\nu_{l}b \quad \text{and} \quad \bar{t} \to W^{-}\bar{b} \to q\bar{q}'\bar{b}$$

Use b quark tag to get very clean $W + b+$ jets sample

- The $W + 1$ and $W + 2$ jet samples are control samples
  - Don't expect significant top contribution

Sample	SVX bkg	SVX tags	SLT bkg	SLT tags
W+1 jet	50 ± 12	40	159 ± 25	163
W+2 jet	21 ± 7	34	46 ± 7	55
W+$\geq$ 3 jet	6.7 ± 2.1	27	15.4 ± 2.3	23

- Probability of background fluctuation is 1×10^{-6}
  - Combining SVX, SLT and DIL channels
- Agrees well with expected rate from $t\bar{t}$ production



$\boxed{\text{Measurement of Top Mass}}$

We use event kinematics to estimate top quark mass

- Require events to have a fourth jet
  - Require $E_T > 8$ and $|\eta| < 2.4$ for 4rth jet
  - High efficiency ($\sim 85\%$ for $t\bar{t}$)
- Combine all possible jet pairs to form $W \to q\bar{q}$
  - Choose one with best χ^2
  - Require $\chi^2 < 10$
- Use b-tag information to identify b quark jet
- Fit events requiring kinematic constraints

This selection leaves us with 19 tagged events

- Estimate background to be $6.9^{+2.5}_{-1.9}$ events
  - Primarily QCD W+jet

Result of Mass Fits

The fitted mass for the 19 events is consistent with $t\bar{t}$

- Probability that background would fluctuate to give peak with similar significance is $< 2 \times 10^{-2}$



$$M_{top} = 176 \pm 8 \pm 10 \quad \text{GeV/c}^2$$

How Does It All Fit in the SM?

Interesting that everything now points to heavy top

W Mass vs. Top Mass



CDF Long-Range Plan

We have short run in fall (Run 1C)

- Will increase Run 1 data sample to $\sim 150\ \mathrm{pb}^{-1}$
- End in Feburary 1996

Collider programme starts up in early 1999

- Expect to achieve luminosities of $2 \times 10^{32}\ \mathrm{cm}^{-2}\mathrm{s}^{-1}$
- Bunch spacing reduces to 396 and then 132 ns
- 2-year run will collect $\sim 2\ \mathrm{fb}^{-1}$

Collaboration has to upgrade detector over next 4 years

- New plug/forward calorimeter (SciFi)
- SVX II: 5-layer double-sided detector
- Intermediate Fibre Tracker (IFT): 6 layers of SciFi
- Extension of muon coverage (complete azimuthal coverage and move FMU system closer)
- New electronics, Trigger, DAQ, Online and Offline systems
- New central tracking detector

Top Physics Impact

Biggest impact Run 2 will have is increased statistics

- $\times 5 - 10$ in integrated luminosity
- More complete b tagging

Top yields can be reliably estimated from Run 1 data

Channel	1 fb^{-1}	10 fb^{-1}
Tagged $W+\geq 4$ jets	600	6000
Double tagged $W+\geq 4$ jets	300	3000
Tagged Dilepton events	100	1000
$Z+\geq 4$ jet events	200	2000

Most measurements will scale with sensitivity

- $\sigma_{t\bar{t}}$ uncertainty $\pm 5\,(\mathrm{stat}) \pm 7\,(\mathrm{syst})\%$
  - Currently theoretical uncertainty is $\sim \pm 20\%$
- M_{top} measurement is systematics limited

$$\delta(M_{\mathrm{top}}) \sim 3\ \mathrm{GeV/c}^2$$

How Well Can We Measure M_{top}?

Top mass statistical uncertainty will go like $1/\sqrt{\mathcal{L}}$

- Will go down to ~ 2 GeV/c^2 for Run II
- Reduce by using dileptons or multijet events?

Systematic uncertainty driven by jet reconstruction

- Best control is $W \to q\bar{q}'$ signal
- Also have $Z + 1$ and $\gamma + 1$ jet as one calibration
- Expect to reduce $\sigma_{\rm jet}$ to ~ 5 GeV/c^2 in Run I
  - Use W signal and higher statistics control samples
- Should scale as $1/\sqrt{\mathcal{L}}$
  - Experience of W mass instructive
  - Should be able to get to $\sim 1.5 - 2.0$ GeV/c^2

Other uncertainties (6.4 GeV/c^2) should scale

- knowledge of background shape and b tagging bias
- These should reduce to ~ 1.7 GeV/c^2
- Gives total systemaatic uncertainty of 2.4 GeV/c^2

$$\delta(M_{\rm top}) \sim 3 \text{ GeV/c}^2$$

Run II M_{top} and M_W measurements will constrain SM



$\boxed{\textbf{How About } V_{tb}\textbf{?}}$

Limits on V_{tb} depend on V_{ts} and V_{td}

- Best to quote uncertainty on $BR(t \to W^+ b)$
- Will remain dominated by statistical uncertainties
  - Biggest systematic uncertainty is tagging efficiency
  - Will have $\sim 10^7$ $B \to lX$ events to measure this

Uncertainty from $W+$ jets measurement is $\sim \pm 20\%$

- Should get to statistical uncertainty of $\sim \pm 3\%$
- Dileptons will measure BR to $\sim \pm 4\%$
- Should continue to scale with $1/\sqrt{\mathcal{L}}$

With SM constraints on V_{ts} and V_{td}

- Set limit of $V_{tb} > 0.25$ at 95% CL