Top to Bottom

New Results on Top and Bottom Quark Decays

Pekka K. Sinervo
University of Toronto

1. Top and Bottom Quark Decays in the SM
2. Summary of Run I Data Samples
3. Top Quark Branching Fractions
4. Bottom Meson Hadronic Decays
5. Summary and Outlook for Run II

HEP Seminar
October 1996

SM Top Quark Phenomenology

Top quarks decay via the weak force

- Predict $\sim$ 100% $t \to W^+ b$
- Lifetime depends on mass
- For heavy top ($M_{top} \gtrsim 100 - 120$ GeV/c^2), decays before it hadronises

Can be viewed as a "pure" charged current

$$\begin{aligned} \Gamma_{\text{top}} &= \frac{G_F M_t^3}{8\sqrt{2}\pi} \left(1 - \frac{M_W^2}{M_t^2}\right)^2 \left(1 + 2\frac{M_W^2}{M_t^2}\right) \\ &\sim 175 \text{ MeV} \left(\frac{M_t}{M_W}\right)^3 \sim 2 \text{ GeV} \end{aligned}$$

- Toponium doesn't have time to form
- T mesons $\sim$ degenerate (if there was time...)

Decay dominated by "longitudinal W^+"

- Means the W^+ prefers helicity 0
- Result of large top and W^+ mass

What Do Top Decays Tell Us?

SM predicts "pure" $V-A$ decay

- clear test of polarisation at decay vertex

$$\frac{W_{\text{long}}}{W_{\text{T}}} \sim \frac{1}{2}\left(\frac{M_t}{M_W}\right)^2 \sim 2.2$$

- V_{ts} CKM matrix elements
- Constraints on other SM parameters (eg. A_{LR})
- Effectively another consistency check of SM

Sensitive to some classes of "new physics"

- Higgs couples strongly to top
- Could decay into a SUSY H^+

$$t \rightarrow H^+ b$$
$$\hookrightarrow \tau^+ \nu_\tau$$

- Top quark might be "special" in unexpected way
  - New physics that couples to $t\bar{t}$ system

SM Bottom Quark Decays

SM predicts similar charged-current decays

- Dominated by weak matrix element
- However, b is imbedded in "brown muck"
  - Strong interaction effects are important
  - higher-order diagrams also play a role

Have resorted to an effective Hamiltonian approach

- Use a basis convenient for a $Q\bar{q}$ system
  $\Rightarrow$ Heavy Quark Effective Theory
- Assume that decay amplitude "factors" (BSW)
- Determine parameters empirically

Hadronic Bottom Decays

Start with

$$H = \frac{G_F}{\sqrt{2}} V_{cb} \left\{ \left[(\bar{d}u) + (\bar{s}c) \right] (\bar{c}b) \right\}$$

- Renormalise H to take into account QCD corrections

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} \Big\{ c_1(\mu) \left[(\bar{d}u) + (\bar{s}c) \right] (\bar{c}b) + c_2(\mu) \left[(\bar{c}u)(\bar{d}b) + (\bar{c}c)(\bar{c}b) \right] \Big\} \quad (1)$$

- For $B \to J/\psi$ decays, employ BSW factorisation:

$$d\Gamma(B^+ \to J/\psi X) \propto a_2(m_b)^2 |V_{cb} V_{cs}^*|^2 \, |\langle J/\psi | (\bar{c}c) | 0 \rangle|^2 \, |\langle X | (\bar{s}b) | B \rangle|^2 \quad (2)$$

- Have to put in b quark form factors by hand

Model is fundamental enough to warrant detailed study

- CLEO/ARGUS have looked at many decays systematically
- But studies have had limitations
  - low statistics
  - only produce B° & B^+ (no B_s° or baryons)

Hadron collider experiment overcomes some limitations

- Tevatron has produced $\sim 3 \times 10^9$ $b\bar{b}$ events

Run I Results from CDF

CDF is general-purpose detector designed to study $\overline{p}\,p$ collisions

- approximately hermetic calorimeters in $|\eta| < 4.2$
- 84-layer drift chamber in 1.4 T solenoidal B field
- muon chambers outside calorimeter
- electron ID based on shower profiles & tracking

The CDF experiment ran from 1992-96

- Collected 20 pb^{-1} in 1992-93 (Run Ia)
- Collected 85 pb^{-1} in 1994-96 (Run Ib)
- Maximum L of $\sim 2.7 \times 10^{31}$ $\text{cm}^{-2}\text{s}^{-1}$

Top Quark Production

Top quark production is mediated by

$$q\bar{q}, gg \quad \rightarrow \quad t\bar{t}$$

$q\bar{q} \rightarrow t\bar{b}X$ and $q\bar{q} \rightarrow W^+ \rightarrow t\bar{b}$ both small

Because top is heavy, need large x for initial partons

- Implies that cross section falls with mass
- $t\,\bar{t}$ system approximately at rest – $< P_T > \sim M_{top}/2$

Higher order corrections to $\sigma_{t\,\bar{t}}$ have been calculated

- Most recently by Laenen, Smith and Neerven, NP **B369**, (1992) 543.
- Biggest effects due to "soft gluon emission"



Top Decays and Detection Requirements

SM top quark decays via the weak force

$$t \rightarrow W^+ b$$
$$\hookrightarrow u\bar{d} \text{ or } c\bar{s} \quad (2/3)$$
$$\hookrightarrow e^+\nu_e \text{ or } \mu^+\nu_\mu \text{ or } \tau^+\nu_\tau \quad (1/3)$$

In proton-antiproton collisions,



This results in several distinct decay signatures

- Six jet final states (36/81)
- Lepton ($e^\pm$ or $\mu^\pm$) + Jet events (24/81)
- Dilepton events with 2 energetic $e^\pm$ and $\mu^\mp$ (4/81)
- Decay modes with τ's hard! ($\sim$ 12/81)

Triggering Issues

Use multi-level trigger to tag events in real-time

- Interaction rate of

  $\left(50 \times 10^{-27}\ \text{cm}^2\right) \times \left(1.5 \times 10^{31}\ \text{cm}^{-2}\text{s}^{-1}\right) = 7.5 \times 10^5$ Hz

  has to be reduced to of order 1-10 Hz

- Most challenging aspect of experiment
  - Depends critically on all subsystems
  - Need efficiency $> 75\%$ to be competitive

Triggering for "heavy" top is relatively easy

- Tag high P_T ($> 15 - 20$ GeV/c) lepton(s)
- Missing energy requirements ($\not\!\!E_T >$ $20 - 25$ GeV)
- Multijet topologies
- Trigger efficiencies $\geq 90 - 95\%$ in most cases

Top Signals

Have detected signals for $t\bar{t}$ production & decay

1. Lepton + jets channel
   - Require $e^{\pm}$ or $\mu^{\pm}$ with $P_T > 20$ GeV/c
   - Require $\not\!\!E_T > 20$ GeV to tag neutrino
   - Require ≥ 3 jets with $E_T > 15$ GeV
   - Require at least one jet with b tag
2. Dilepton channel
   - Require two $e^{\pm}$ or $\mu^{\pm}$ with $P_T > 20$ GeV/c
   - Require $\not\!\!E_T > 25$ GeV to tag neutrino
   - Require ≥ 2 jets with $E_T > 10$ GeV
3. Dileptons with τ's
   - Require τ candidate + additional lepton
   - Similar requirements to standard dileptons
4. Hadronic channel
   - Require 6 jets with $E_T > 20$ GeV
   - Require at least two jets with b tags
   - Make kinematic cuts to increase S/N

W+Jet Selection

Selection of W candidates straightforward

- Require high P_T isolated lepton candidate
- Require significant $\not\!\!E_T$
- Require three or more high P_T jets

Have large W^+ samples

- CDF has 31,000 $e\nu_e$ and 21,000 $\mu\nu_\mu$ candidates

Run IA CDF data shows clear "W" signal:



W + Jet Sample

Require $\geq$ 3 or 4 jets (from other W and b quarks)

- $\geq$ 3 jets with $E_T > 15$ GeV and $|\eta| < 2.0$
- find 274 W+$\geq$ 3 jet events

Most background sources are small:

Background	Fraction of Events (%)
Fake leptons and $b\bar{b}$	9 ± 5
WW, WZ and ZZ	4 ± 2
$Z \rightarrow \tau\tau$, ee, $\mu\mu$	7 ± 2
Total	20 ± 6

QCD W+jets production is largest expected background

M_{top} (GeV/c^2)	Number of Events (CDF)
120	176 ± 28
140	105 ± 14
160	57 ± 10
180	16 ± 3.2
W+jets	238^{+76}_{-38}

b Quark Tagging in W+Jet Events

Have two different techniques for b-tagging

1. About 20% of b's decay via

$$b \to e^- \nu_e X \quad \text{and} \quad b \to \mu^- \nu_\mu X$$

   - Look for jets with "soft" leptons in them (SLT tags)

2. "Long" b lifetime – $c\tau \sim 450$ microns

   - Use Silicon Vertex Detector to identify jets with a secondary vertex
   - Expect 42% of $t\bar{t}$ events to have at least one SVX tag

Applied b tagging techniques to W+jet events

- The $W + 1$ and $W + 2$ jet samples are ultimate control
  - Don't expect significant top contribution
  - Theory predicts low b hadron content

Sample	SVX bkg	SVX tags	SLT bkg	SLT tags
W+1 jet	69 ± 15	70	272 ± 24	245
W+2 jet	29 ± 4	45	78 ± 7	82
W+$\geq$ 3 jet	9.3 ± 1.1	34	25.2 ± 3.0	40

Dilepton Top Sample

$t\bar{t}$ decaying into 2 leptons create unique signature

- Two charged leptons from $W^{\pm}$ decay
- Imbalance in total energy due to undetected neutrinos
- Two jets from bottom quarks

Select sample by requiring

- Two clean, high P_T lepton candidates
- Missing transverse energy $\not\!\!E_T > 25$ GeV
- Require two jets with $E_T >$ 10 GeV and $|\eta| < 2.0$
- Reject specific backgrounds (e.g. Z°)

Run 1 ee data (109 pb^{-1}), CDF preliminary



Dilepton Channel

Significant dilepton background is Drell-Yan production

- $\not\!\!E_T$ signature created by mismeasured jet
- Harden $\not\!\!E_T$ cut to > 50 GeV if correlated with jet



a) Z+1 jet, b) Z+ 2 jet, and c) $t\bar{t}$ ($M_{\text{top}} = 160$ GeV/c^2)

Dielectron Data for Run 1A and 1B

Run 1 ee data (109 pb^{-1}), CDF preliminary



DiMuon Data for Run 1A and 1B

Run 1 μμ data (109 pb^{-1}), CDF preliminary



Electron-Muon Data for Run 1A and 1B

Run 1 eμ data (109 pb^{-1}), CDF preliminary



Summary of CDF Dilepton Search

Observe a total of 10 candidate events

- Observe 1 e^+e^- event, 2 $\mu^+\mu^-$ event and 7 $e\mu$ events
  - One $\mu^+\mu^-$ event suspicious (but we keep it)
  - Has $\mu^+\mu^-\gamma$ mass close to Z°
- Expect 2.1 ± 0.4 events from background

Acceptance $\times$ SM branching fraction $= 0.77 \pm 0.08\%$

$$\sigma_{t\bar{t}} = 9.3^{+4.4}_{-3.4} \text{ pb}$$



Top Quark Mass from Lepton + Jets

In this mode, have measured almost all $t\bar{t}$ daughters

- Lepton 4-momentum solid and ν_l is missing only p_z
- $W \to q\bar{q}'$ observed as 2 jets
- bottom quarks observed as jets (perhaps tagged)



Problems:

- QCD radiation generates additional jets
- Combinatorics in identifying jets $\leftrightarrow$ partons
- Background events

Top Mass Results

Estimate mass based on 34 $W+ \geq 4$ jet sample

- Use known fake tag rates and $Wb\bar{b}$
- Background shape in M_{top} comes from MC calculation
- Estimate background to be $6.4^{+2.1}_{-1.4}$ events
  - Primarily QCD W+jet



Evidence for W → qq̄

Observation of $W \rightarrow q\bar{q}$ is important

- Strong evidence of signal in $W^+W^-b\bar{b}$ channel
- Can calibrate jet energy scale

Have to refit data relaxing W mass constraint

- Look for W signal plotting all mass combinations
  - Makes it tough since 6 combos/event
- Reduce backgrounds by requiring $H > 300$ GeV



Top Quark Branching Fractions

Indication of new physics would be if $\mathcal{B}(t \to W^{\pm} b) \neq 1$

- SM expects V_{tb} to be unity
  - Unitarity constraints
    - V_{ud} large from nuclear β and μ decay
    - V_{us} constrained by K_{e3} decays
    - V_{cd} comes from $\nu n \to cX$
    - V_{cs} constrained by D_{e3} decays
    - V_{cb} small because large τ_b
    - V_{ub} even smaller ($b \to u$)
  - Expect V_{ts} and V_{td} much smaller than V_{tb}
- $V_{tb} \lesssim 0.9985$ violates (or bends) SM

Experimentally, we determine

$$\mathcal{R} \equiv \frac{\mathcal{B}(t \to Wb)}{\mathcal{B}(t \to Wq)} = \frac{|V_{tb}|^2}{|V_{tb}|^2 + |V_{ts}|^2 + |V_{td}|^2}$$

Measure the rate of 0, 1 and 2 b-tagged events

- Need to be able to b tag
- Have to deal with large backgrounds in 0-tagged sample

$\mathcal{B}$ Technique

Fraction of zero, single and double-tagged events is

$$\begin{aligned} f_0 &= (1-\mathcal{R}\epsilon)^2 \\ f_1 &= 2\mathcal{R}\epsilon(1-\mathcal{R}\epsilon) \\ f_2 &= (\mathcal{R}\epsilon)^2 \end{aligned}$$

- ϵ is tagging efficiency

Rate of single vs double tags gives

$$\mathcal{R} = \frac{2}{\epsilon(f_1/f_2+2)}$$

Rate of zero vs single tags gives

$$\mathcal{R} = \frac{1}{\epsilon(2f_0/f_1+1)}$$

Estimate $\mathcal{R}$ using maximum likelihood technique

- For a given number of observed events,
  - calculate probability of observing 0, 1 and 2 tags
  - maximize probability by varying $\mathcal{R}$
- Can use probability to constraint $\mathcal{R}$

Event Selection Details

Increase sensitivity by relaxing b tag requirement

- Reject large QCD background by cutting on
  1. Relative likelihood of $t\bar{t}$ vs W+jets
     - based on 2nd and 3rd leading jet E_T
     - Rejects 75% of W+jet background
     - Has efficiency of 75% for $t\bar{t}$
  2. Use sample with $|\cos\theta^*| < 0.7$ as check
     - Expected to be enriched in $t\bar{t}$
     - But fairly large systematic uncertainty
- Only using Run Ib data (~ 90 pb^{-1})

After applying likelihood cut,

- Have 68 events after relative likelihood cut
  - Expect non-$t\bar{t}$ background to be ~ 40 events
  - Have 24 events after additional $\cos\theta^*$ cut
- Have 10 and 5 single and double-tagged events

CDF BR Results

For dilepton events

- Have 5, 2 and 0 events with 0, 1 and 2 b-tags
- Backgrounds are very small ($\lesssim 2$ events)

Fit $\mathcal{B}(t \to W^+ b)$ using observed event rates

Class of Events	Observed	Fitted
Lepton + Jets – 0 tags	51.5	53
Lepton + Jets – 1 tags	10.4	10
Lepton + Jets – 2 tags	4.9	5
Dileptons – 0 tags	4.9	5
Dileptons – 1 tags	1.4	2
Dileptons – 2 tags	0.2	0

Fit results are:

$$\mathcal{R} = 1.42^{+0.41}_{-0.39} \quad \text{(lepton + jets)}$$

$$\mathcal{R} = 0.72^{+0.66}_{-0.44} \quad \text{(dileptons)}$$

$$\mathcal{R} = 1.23^{+0.38}_{-0.33} \quad \text{(combined)}$$

- Values > 1 allowed because of background fluctuations

Relating Back to V_{tb}



Find that $\mathcal{R} > 0.61$ at 95% C.L.

- Can relate that to V_{tb} using average values of
  - $0.004 \leq |V_{td}| \leq 0.015$ and $0.030 \leq |V_{ts}| \leq 0.048$

$V_{tb} > 0.78$ (95%C.L.) assuming CKM unitary
$V_{tb} > 0.38$ (95%C.L.) without CKM unitary

- Really saying that $|V_{tb}| \gg |V_{ts}|$ or $|V_{td}|$

Other Features of Top Decays

We have also looked for "exotic" decay modes

- $t \rightarrow Z^{\circ} q$
  - Just run out of statistics
  - Very weak limit on $\mathcal{B}(t \rightarrow Z^{\circ} q)$

Have also looked at polarisation of top decay

- May be just beyond reach statistically

Have yet to perform a complete study using all modes

- Incorporate hadronic modes
- Add in τ decay modes

Bottom Meson Decays at CDF

Hadron collider b experiment has serious S/N problem

- b cross section is of order $\sim$ 30 μb
- light quark and gluon jet $\sigma \sim$ 5 mb
- solution is to trigger on "unique" topologies
  - require semileptonic decays
  - require tag such as $J/\psi \to \mu^+\mu^-$ in final state
- All CDF studies have been based on these triggers

However, can extract clean samples efficiently

- Will focus on $B \to J/\psi X$ and $B \to \psi(2S)X$ decays
- Now have world's largest sample

Why are J/ψ and ψ(2S) States Interesting?

Expect decays to result from "internal" diagrams

- Theoretically expected to be very clean
- Strong test of factorisation
  - Technically considered type 2 decays
  - Only depend on a_2 in BSW model
- Test details of calculations

There has also been some controversy

- Polarisation in $B \to J/\psi K^*(890)$ decay larger than predicted
- Hard to reconcile this with branching fractions
- Few ways out
  - Measurements are wrong!
  - Factorisation only approximate
  - Form factors are just wrong

Measurement Technique

The observed cross section (*e.g.* $B^+ \to J/\psi K^+$) is

$$\sigma_{\text{obs}} = \sigma(p\bar{p} \to \bar{b}) f_u \mathcal{B}(B^+ \to J/\psi K^+) \epsilon^{K^+}$$

There are large uncertainties in $\sigma(p\bar{p} \to \bar{b})$

- Also significant systematic uncertainties in ϵ^{K^+}
- Strategy is to measure ratios of σ_{obs}

$$R^{K^\circ}_{K^+} = \frac{f_d}{f_u} \frac{\mathcal{B}(B^\circ \to J/\psi K^\circ)}{\mathcal{B}(B^+ \to J/\psi K^+)}$$

- We also get ratios of f_i for "free"
  - Have to use independent data or theory
  - However, tests "independent" fragmentation ansatzs

Selection of Exclusive B $\rightarrow$ J/ψX Decays

Search for exclusive decays

$$\begin{aligned} B^{+} &\rightarrow J/\psi K^{+}, \\ B^{\circ} &\rightarrow J/\psi K^{\circ}, \\ B^{+} &\rightarrow J/\psi K^{*}(892)^{+}, \\ B^{\circ} &\rightarrow J/\psi K^{*}(892)^{0}, \\ B_{s}^{\circ} &\rightarrow J/\psi \phi(1020) \end{aligned}$$

Select decays by requiring

- Two muon candidates with $P_T \gtrsim 3$ GeV/c
- Require muons to come from common vertex
- Search combinatorially for daughter mesons
- Require B candidate $P_T > 6$ or 8 GeV/c
- Require meson daughter $P_T > 1.5$ or 2.0 GeV/c
- Require decay length of B candidate to be > 0

J/ψ Candidates

CDF dimuon trigger upgraded in Run I for J/ψ's

- Now extends over $|\eta| \lesssim 1.0$
- Added steel and chambers to reduce backgrounds

$\Rightarrow$ efficient trigger for J/ψ $P_T \gtrsim 6$ GeV/c



Run IA Results











Ratios of Branching Fractions

$$R^{K^\circ}_{K^+} = \frac{f_d}{f_u}\,\frac{\mathcal{B}(B^\circ \to J/\psi K^\circ)}{\mathcal{B}(B^+ \to J/\psi K^+)} = 1.15 \pm 0.27 \pm 0.09,$$

$$R^{K^{*+}}_{K^+} = \frac{\mathcal{B}(B^+ \to J/\psi K^{*+})}{\mathcal{B}(B^+ \to J/\psi K^+)} = 1.92 \pm 0.60 \pm 0.17,$$

$$R^{K^{*\circ}}_{K^+} = \frac{f_d}{f_u}\,\frac{\mathcal{B}(B^\circ \to J/\psi K^{*\circ})}{\mathcal{B}(B^+ \to J/\psi K^+)} = 1.59 \pm 0.33 \pm 0.12,$$

$$R^{K^\phi}_{K^+} = \frac{f_s}{f_u}\,\frac{\mathcal{B}(B^\circ_s \to J/\psi \phi)}{\mathcal{B}(B^+ \to J/\psi K^+)} = 0.41 \pm 0.12 \pm 0.04,$$

$$R^{K^{*+}}_{K^\circ} = \frac{f_u}{f_d}\,\frac{\mathcal{B}(B^+ \to J/\psi K^{*+})}{\mathcal{B}(B^\circ \to J/\psi K^\circ)} = 1.68 \pm 0.58 \pm 0.11,$$

$$R^{K^{*\circ}}_{K^\circ} = \frac{\mathcal{B}(B^\circ \to J/\psi K^{*\circ})}{\mathcal{B}(B^\circ \to J/\psi K^\circ)} = 1.39 \pm 0.36 \pm 0.10,$$

$$R^{K^\phi}_{K^\circ} = \frac{f_s}{f_d}\,\frac{\mathcal{B}(B^\circ_s \to J/\psi \phi)}{\mathcal{B}(B^\circ \to J/\psi K^\circ)} = 0.35 \pm 0.12 \pm 0.03,$$

$$R^{K^{*\circ}}_{K^{*+}} = \frac{f_d}{f_u}\,\frac{\mathcal{B}(B^\circ \to J/\psi K^{*\circ})}{\mathcal{B}(B^+ \to J/\psi K^{*+})} = 0.83 \pm 0.27 \pm 0.07,$$

$$R^{K^\phi}_{K^{*+}} = \frac{f_s}{f_u}\,\frac{\mathcal{B}(B^\circ_s \to J/\psi \phi)}{\mathcal{B}(B^+ \to J/\psi K^{*+})} = 0.21 \pm 0.08 \pm 0.02,$$

$$R^{K^\phi}_{K^{*\circ}} = \frac{f_s}{f_d}\,\frac{\mathcal{B}(B^\circ_s \to J/\psi \phi)}{\mathcal{B}(B^\circ \to J/\psi K^{*\circ})} = 0.26 \pm 0.08 \pm 0.02.$$

Comparison with HQET and BSW

Have compared results with several model predictions

- A. Deandrea, N. Di Bartolomeo, R. Gatto and G. Nardulli, Phys. Lett. B **318**, 549 (1993).
- Other calculations give similar results



Fragmentation Fractions

Can extract fragmentation fractions

- Use Deandra *et al.* model predictions
- Combine measurements from different channels

$$\frac{f_d}{fu} = 0.99 \pm 0.19 \pm 0.08 \quad \text{and} \quad \frac{f_s}{(fu, f_d)} = 0.34 \pm 0.10 \pm 0.03.$$

Can estimate absolute fragmentation fractions

- Assume that $f_u + f_d + f_s + f_{\Lambda_b} = 1$
- Assume that $f_{\Lambda_b} \sim 0.115$

$$f_u = \frac{1 - f_{\Lambda_b}}{1 + \frac{f_d}{f_u} + \frac{f_s}{f_u}}$$

- This gives us

$$\begin{aligned} f_u &= 0.39 \pm 0.04 \pm 0.04, \\ f_d &= 0.38 \pm 0.04 \pm 0.04 \text{ and} \\ f_s &= 0.13 \pm 0.03 \pm 0.01. \end{aligned}$$

In good agreement with PDG "LEP" values

- $f_u : f_d : f_s = 0.378 \pm 0.022 : 0.378 \pm 0.022 : 0.112 \pm 0.019$

$\psi(2S)$ Branching Fractions

Currently working on measuring ratios

$$\frac{\mathcal{B}(B^+ \to \psi(2S)K^+)}{\mathcal{B}(B^+ \to J/\psi K^+)}$$
$$\frac{\mathcal{B}(B^0 \to \psi(2S)K^{*0})}{\mathcal{B}(B^0 \to J/\psi K^{*0})}$$

- Using full Run I data sample
- See clean inclusive $\psi(2S)$ signal

$$\psi(2S) \to J/\psi\pi^+\pi^- \text{ and } \psi(2S) \to \mu^+\mu^-$$





Run IB $B \to \psi(2S)X$ Signals











Preliminary Results

The ratios of branching fractions are

$$\frac{\mathcal{B}(B^+ \to \psi(2S)K^+)}{\mathcal{B}(B^+ \to J/\psi K^+)} = 0.666 \pm 0.093 \pm 0.101$$
$$\frac{\mathcal{B}(B^0 \to \psi(2S)K^{*0})}{\mathcal{B}(B^0 \to J/\psi K^{*0})} = 0.569 \pm 0.131 \pm 0.074$$

Can turn these into absolute branching fractions using

$$\mathcal{B}(B^+ \to J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$$
$$\mathcal{B}(B^0 \to J/\psi K^{*0}) = (1.58 \pm 0.28) \times 10^{-3}$$

This gives us the branching fractions

$$\mathcal{B}(B^+ \to \psi(2S)K^+) = (6.8 \pm 1.0 \pm 1.4) \times 10^{-4}$$
$$\mathcal{B}(B^0 \to \psi(2S)K^{*0}) = (9.0 \pm 2.1 \pm 2.0) \times 10^{-4}$$

Comparison with Other Measurements

These results can be compared with CLEO results



Now the world's best measurement

Summary

So far, top decays just like the SM predicts

- Constraints on $V_{tb} > 0.38$ at 95% C.L.
- Polarisation studies underway
- Nothing anomolous found

Bottom decay studies really only starting

- Expect to have the most precise $B \to J/\psi$ and $B \to \psi(2S)$ measurements in next year
- Polarisation studies of entire Run I sample underway
- Will allow tests of HQET and BSW factorisation

Run II expected to start in 1999

- Integrated luminosity of ~ 2 fb^{-1}
- Upgraded tracking, calorimetry and DAQ
- Will increase top and bottom samples by $\times 50$